



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>







HELP TO ARITHMETIC.

Cambridge:

PRINTED BY C. J. CLAY, M.A.
AT THE UNIVERSITY PRESS.

HELP TO ARITHMETIC

DESIGNED FOR THE USE OF
SCHOOLS.

BY

H. CANDLER, M.A.,

OF TRINITY COLLEGE, CAMBRIDGE;
AND MATHEMATICAL MASTER OF UPPINGHAM SCHOOL.



London and Cambridge:
MACMILLAN AND CO.
1868

[All Rights reserved.]

181. 9 37.

PREFACE.

THE writer of the following pages believes that an apology is due for the apparent want of completeness and consecutiveness that are evident in his treatment of the subject they deal with. His only excuse is that this very inconsecutiveness was intentional. His desire was not to make a book. That would seem unnecessary in the presence of the many very excellent treatises on Arithmetic now current in schools. The present work is intended as a companion to any text-book that may be in use, and the object of the writer has been twofold. He has proposed to himself to exhibit in a compendious form explanations of certain Arithmetical difficulties, which, in the course of many years' teaching, he has found satisfactory to his pupils; and to put before boys examples of sums in various rules fully worked out in a clear, uniform, and natural manner. In the "money rules" he has endeavoured

to show that there are no new processes ; only new technical expressions to be explained ; and that these rules can therefore be made at once to depend on the ordinary forms of proportion.

It will be seen from the above remarks that the design of the book is of the humblest nature. There are no novelties, no ambitious attempts. It only aspires to be useful, and endeavours to accomplish this end by means of clear explanations, and by urging upon learners the necessity of a neat, logical, and methodical treatment of sums.

It is the intention of the writer to follow up this publication with two or more similar treatises on Algebra, Euclid, and other kindred school subjects of an elementary nature ; the whole will thus form a single work, of which the present must be considered the First Part.

CONTENTS.

	PAGE
I. INTRODUCTION	1
Numeration and Notation	2
Division, Definition of	2
Arithmetical Tables	3
Square and Cubic Measure	5
Least Common Multiple	9
II. Vulgar Fractions	11
III. Decimal Fractions	28
IV. Proportion	40
V. RULES REDUCIBLE TO PROPORTION.	
Simple Interest	43
Compound Interest	47
Discount	50
Stocks	51
Profit and Loss	56
VI. MISCELLANEOUS EXAMPLES.	
Proportional Parts, &c.	59
Chain Rule	62
Per centage	63
Concluding Remarks	63
NOTES	68

PART I.

ARITHMETIC.

1. IN order to avoid using a new symbol and word for every different number, all nations, who have attained any proficiency in counting at all, have adopted a radix, or turning point, after which no new symbols are used ; and farther numbers are expressed by the combination of old symbols, and by the positions which they relatively occupy. This radix has in all civilized nations been the number ten ; not that that is the most convenient number, (twelve would be the most convenient), but because the system was adopted while they were still in a savage state, and did not reason on its convenience or inconvenience for elaborate computation. They were led to employ the number ten by reason of the ten fingers of the hand, and no doubt at first counted with small round stones or any similar substances. The words *digit* and *calculation* refer us back to a rude origin of Arithmetic, or rather, of counting. A clear and interesting explanation of numeration and notation will be found in the early part of De Morgan's Arithmetic.

2. Numeration and Notation.

Begin at the unit figure and separate the figures into groups of six each. Then the figures in the second group will give the number of millions, in the third group the number of billions, in the fourth group the number of trillions, and so on.

Thus, take the number

12345678905698.

Group it thus :

12,345678,905698.

It will be read as follows :

12 billions, 345 thousand 678 millions, 905 thousand, 698.

N.B. The rule given above is not universally adopted. Some writers, (all continental nations we believe), consider a billion to be a thousand millions, a trillion, a thousand billions, and so on. In this way, the above number would be read, 12 trillions, 345 billions, 678 millions, 905 thousand, 698.

3. Division.

The definition of division sometimes given in Arithmetical treatises, is, that it is "the process of finding how often one number is contained in another." This scarcely seems to be sufficient, for it does not include all classes of Arithmetical cases. For abstract numbers it is true, but not necessarily for concrete numbers. In order to take in all cases, and to retain strictly Arithmetical language and ideas, it would seem that two definitions are necessary, as thus.

By the division of one concrete quantity by another *of the same kind* is meant the process of finding how

often the second is contained in the first, or how many parcels, each containing the second number of things, there will be in the first number of things: e. g. 12 apples \div 3 apples. Ans. 4.

By the division of a concrete quantity by an abstract number, is meant the process by which we find, if we separate the first into as many equal parcels as are represented by the second, how many units each parcel will contain: e. g. £2. 16s. \div 3. Ans. 18s. 8d.

We shall return to these definitions in Note B, on the Division of Fractions.

4. *Arithmetical Tables.*

The following tabular results are very important, and are far too often forgotten.

In Avoirdupois weight ;

112 lbs. make 1 cwt.

The lb. Av. contains 7000 grs. Tr.

In Length measure ;

6 ft. make 1 fathom.

$5\frac{1}{2}$ yds. make 1 rod, pole, or perch.

220 yds. make 1 furlong.

1760 yds. make 1 mile.

In Cloth measure ;

4 qrs. make 1 yd.

5 qrs. make 1 English ell.

In measure of Surface ;

$30\frac{1}{4}$ sq. yds. make 1 sq. rod, pole, or perch.

4840 sq. yds. make 1 acre.

In Cubic measure ;

1728 cubic in. make 1 cubic ft.

The following weights and measures, though not required for ordinary Arithmetic, should be observed.

In Length measure ;

12 lines make 1 inch.

9 inches make 1 span.

18 inches make 1 cubit.

$69\frac{1}{9}$ miles make 1 geographical degree.

$69\frac{1}{9}$ miles is the length of $\frac{1}{360}$ part of the circumference of the earth at the equator.

In Cloth measure ;

A Flemish ell contains 3 qrs.

A French ell contains 6 qrs.

In Capacity ;

A firkin (of beer) contains 9 gals.

A kilderkin “ contains 18 gals.

A barrel “ contains 36 gals.

A hogshead “ contains 54 gals.

A pipe (of wine) contains 126 gals.

Value of Coins ;

A groat = 4*d*.

A tester = 6*d*.

A noble = 6*s*. 8*d*. $\left(= \frac{\pounds 1}{3} \right)$.

An angel = 10*s*.

A mark = 13*s*. 4*d*. $\left(= \frac{\pounds 2}{3} \right)$.

A moidore = 27*s*.

In Surveying ;

Gunter's chain is 22 yds. long, and contains 100 links.

Hence an acre contains 10 square chains.

A cub. foot of distilled water weighs 1000 oz.

A lb. of gold contains 24 carats.

From this fact we express the fineness of gold by saying that it is so many carats fine ; i.e. so many parts out of 24.

Thus, if a mass of gold were 15 carats fine, $\frac{15}{24}$ of the mass would be gold, and the rest, viz. $\frac{9}{24}$, would be "alloy," a substance depending on the nature of the mass.

Standard gold is 22 carats fine, the remaining two parts being a mixture of silver and copper. Nearly $46\frac{3}{4}$ sovereigns are coined out of a lb. Tr. of standard gold. Jewellers' gold is 18 carats fine.

Standard silver contains 37 parts out of 40 pure silver, the remaining three parts being copper. A lb. Tr. of standard silver yields 66 shillings.

24 pence are coined out of a lb. Av. of copper.

5. *Square and Cubic Measure.*


A common or linear yard is length, and measures length.

A sq. yard is surface, and measures surface, or area. It is a square surface whose length and breadth are both a yard. The length round a sq. yard is 4 yards.

A cubic yard is a solid, and measures solidity, or cubic contents. It is a solid whose length, breadth, and height are each a yard.

A cubic yard has 6 surfaces, each of which is a sq. yard.

From these facts it is clear that the ideas of a common, a square, and a cubic yard, are perfectly different. No amount of common yards will make a square yard. There is a very intimate relation between common and square yards, as there is between apples and pence, but as no amount of pence will make apples, so no amount of common yards will make square yards.



When the length table is known the surface table can be formed from it by multiplying each of the figures of the length table by itself.

This is easy to prove. For let $ABCD$ be a square yard. Then both AB and BC are common yards. Therefore each of them consists of three feet. If through the extremities of these feet we draw straight lines parallel to the sides of the square, it is clear that we shall get 9 compartments each of which is a foot long and a foot broad, i.e. we shall have 9 sq. ft.; which proves the statement.

Thus 12 in. make 1 ft.

And 12×12 sq. in. make 1 sq. ft.

Again, $5\frac{1}{2}$ yds. make 1 p.

And $5\frac{1}{2} \times 5\frac{1}{2}$ sq. yds. $\left(\text{i.e. } \frac{11}{2} \times \frac{11}{2} = \frac{121}{4} = 30\frac{1}{4} \right)$ make one sq. pole.

So in cubic measure, we must multiply any figure of the common table three times by itself to get the corresponding figure of the cubic table.

For along all the lines of the above figure build walls, and, when the walls are a foot high, lay down a floor; then there will be 9 boxes each a foot long, a foot broad and a foot high; that is, there will be 9 cubic feet. Repeat this till our walls are three feet, or a yard high. There then will be three times 9 or 27 cubic feet in the solid. But the solid is a yard long, a yard broad, and a yard high, that is, it is a cubic yard. Hence there are 27 cubic feet in a cubic yard; which proves the statement.

Thus 12 in. make 1 ft.

And $12 \times 12 \times 12$ cubic in. make 1 cubic ft.

It is easy to prove in exactly the same manner as above, that if we want to find the number of sq. ft. in any area, we must multiply the number of feet in the length by the number of feet in the breadth. So we can find the number of cubic feet in a solid by multiplying the number of feet in the length, the number of feet in the breadth, and the number of feet in the height, together.

It must be remembered that length cannot be multiplied by breadth. We are however at liberty to use the expression "length \times breadth = area," if we understand that we are using an abbreviation, in order to avoid the continual repetition of a long phrase. The same is clearly true for cubic measure.

Length cannot be multiplied by breadth, because that is contrary to the definition of multiplication. Nor could length and breadth be said in any arithmetical sense to make area. We shall return to this point in Note A, on the multiplication of fractions.

The following three formulæ are sufficient for all questions in square and cubic measure :—

$$l \times b = a \dots\dots\dots (1),$$

$$l \times b \times h = c \dots\dots\dots (2),$$

$$a \times h = c \dots\dots\dots (3),$$

where either l , b , h , stand for length, breadth or width, height or depth, or thickness ; a stands for area or surface, and c for cubic contents, or solidity.

Ex. There is a room which is 12 feet long, 9 feet broad and 8 feet high. Find

1. The area of the ceiling.
2. The area of the walls.
3. The area of the whole of the interior of the room.
4. The length of the sum of all the edges of the room.
5. The length of a beading round the ceiling.
6. The area of a cornice, three inches broad, in the ceiling round the room.
7. The area of a wainscoating, nine inches broad, in the walls round the floor.
8. The thickness of sand laid evenly over the floor, the cubic dimensions of the sand originally being 8 ft. long, 6 ft. broad and 8 in. deep.
9. The cubic contents of the room.

6. *Least Common Multiple.*

One of two methods must be carefully insisted on. Either the "divisors" must all be *prime*, or care must be taken to divide each number by the largest number that will divide both it and the "divisor" in question. In the latter case the "divisor" must divide one at least of the numbers exactly, in order to *ensure* a correct result.

Ex. L.C.M. of 2, 3, 6, 12, 18, 20, 25, 27, 32, 45.

First Method.

$$\begin{array}{r}
 2 \overline{) 2, 3, 6, 12, 18, 20, 25, 27, 32, 45.} \\
 \quad 2 \overline{) 6, 10, 25, 27, 16, 45.} \\
 \qquad 3 \overline{) 3, 5, 25, 27, 8, 45.} \\
 \qquad \quad 3 \overline{) 25, 9, 8, 15.} \\
 \qquad \qquad \quad 25, 3, 8, 5.
 \end{array}$$

Ans. = $25 \times 3 \times 8 \times 3 \times 3 \times 2 \times 2 = 21600$.

Second method.

$$\begin{array}{r}
 12 \overline{) 2, 3, 6, 12, 18, 20, 25, 27, 32, 45.} \\
 \quad 3 \overline{) 1, 3, 5, 25, 9, 15.} \\
 \qquad \quad 25, 3, 8, 5.
 \end{array}$$

Ans. = $25 \times 3 \times 8 \times 3 \times 12 = 21600$.

Thus in the 2nd line of the 2nd method, under the 18 we put a 3, because the largest number that will divide both 18 and the "divisor" 12, is 6, and 6 will divide 18 with quotient 3. Again the largest number that will divide both 20 and the "divisor" 12 is 4, and 4 will divide 20 with quotient 5: so that we place a 5 under the 20. And so on.

It is clear that the second method is shorter, but requires

but the *whole* quantity. Again if the numerator is greater than the denominator (e.g. $\frac{9}{7}$) it is clear that the expression "the fraction of a quantity" is a confusion in terms (for we could not divide a quantity into 7 parts and take 9 of them). In such a case we must suppose any number of quantities, divide them each into the required number of parts and take the required number of those parts.

From this it is manifest that an improper fraction is equal to (if the numerator = the denominator) or is greater than (if the numerator is greater than the denominator) the whole quantity.

N.B. We have here slightly departed from the language of Arithmetical treatises, which call a fraction whose numerator is equal to its denominator a proper fraction.

Another definition of a fraction as arising from the first, is sometimes added, viz., that it equals the numerator divided by the denominator. We do not think the proof is quite satisfactory without generalizing the meaning of our words in a manner not fitting to pure arithmetic, nor do we think this the right place either to prove or adopt another definition. We shall return to this in the articles on Division of Fractions, merely stating that it will not be necessary for us to use this second definition till then.

8. *To reduce a mixed number to an improper fraction, and conversely.*

$$\text{Ex. } 3\frac{5}{7} = \frac{3 \times 7 + 5}{7} = \frac{21 + 5}{7} = \frac{26}{7}.$$

$$\text{For } 1 (= 7 \text{ sevenths}) = \frac{7}{7}, \text{ therefore } 3 (= 21 \text{ sevenths}) = \frac{21}{7},$$

$$\text{therefore } 3\frac{5}{7} = \frac{21 + 5}{7} = \frac{26}{7}.$$

So on the other hand $\frac{39}{5} = 7\frac{4}{5}$.
$$\begin{array}{r} 5 \overline{) 39} (7 \\ \underline{35} \\ 4 \end{array}$$

For $\frac{5}{5} = 1$, therefore $\frac{35}{5} = 7$, therefore $\frac{39}{5} = 7\frac{4}{5}$.


9. *To multiply a fraction by any whole number or integer.*

Rule. Either multiply the numerator or divide the denominator by it.

Thus $\frac{3}{7} \times 5 = \frac{15}{7}$; $\frac{5}{24} \times 4 = \frac{5}{6}$.

The first rule is clear, for as 3 apples multiplied by 5 is 15 apples, so 3 sevenths ($\frac{3}{7}$) multiplied by 5 is 15 sevenths ($\frac{15}{7}$).

The next rule may be explained thus: we may multiply $\frac{5}{24}$ by 4, either by taking four times as many twenty-fourths, or by taking the 5 parts each four times as large.



But, as we see in the figure, a sixth is four times as large as a twenty-fourth, (for we may divide the 24 parts into six groups of four parts), therefore $\frac{5}{24} \times 4 = \frac{5}{6}$.


10. *To divide a fraction by any integer.*

Rule. Either divide the numerator or multiply the denominator by it.

Thus, $\frac{10}{7} \div 2 = \frac{5}{7}$; $\frac{2}{5} \div 3 = \frac{2}{15}$.

The first rule is clear, for as 10 apples divided by 2 is 5 apples, so 10 sevenths $\left(\frac{10}{7}\right)$ divided by 2 is 5 sevenths $\left(\frac{5}{7}\right)$.

The next rule may be explained thus: we can divide $\frac{2}{5}$ by 3, if we take the two parts each a third of their original size; but a third of a fifth, as we see in the figure,



is a fifteenth, therefore $\frac{2}{5} \div 3 = \frac{2}{15}$.


11. *To reduce a fraction to lower terms.*

Rule. Divide both numerator and denominator by the same number.

Thus, $\frac{6}{15} = \frac{2}{5}$.

The reason of this rule might easily be gathered from the second part of Art. 9. We will however prove it independently.

$\frac{6}{15}$ of a quantity is obtained by dividing it into 15 parts, and taking six of them. But we shall obtain the same result if we take only a third of the number of parts when each of the parts is three times as large. But a third of the number of parts gives us 2 parts. And if the parts are three times as large (see figure) they will be fifths,



for we may divide the 15 parts into 5 groups of 3 parts each. Therefore $\frac{6}{15} = \frac{2}{5}$.

$$\begin{aligned} \text{Ex. } \frac{22176}{23328} &= \frac{11088}{11664} = \frac{5544}{5832} = \frac{2772}{2916} = \frac{1386}{1458} \\ &= \frac{693}{729} = \frac{231}{243} = \frac{77}{81}. \end{aligned}$$

Practical Rule. Divide numerator and denominator by 2 as often as possible, then by 3 as often as possible, similarly by 5, by 7, and by 11. Find the G.C.M. only as a last resort.

Rules are given for finding when a number is divisible by 2, 3, 5, and 11 in Art. 44.

N.B. If the numerator and denominator of a fraction be separated into such factors as can easily be seen by inspection, we may in almost all cases avoid finding the G.C.M. This is perhaps not a good method to teach those who are beginning fractions, but it will be found the shortest means in the end. It is an excellent method of detecting errors and checking work. Numerous rules may be given for accelerating the work of splitting a number into its prime factors.

12. *Division of concrete numbers by integers.*

Ex. 1. Divide £9. 7s. 3½d. by 4.

$$\begin{array}{r} \text{£.} \quad \text{s.} \quad \text{d.} \\ 4 \overline{) 9 \quad 7 \quad 3\frac{1}{2}} \\ \underline{2 \quad 6 \quad 9\frac{1}{2}} \end{array}$$

$$\begin{aligned} 3\frac{1}{4} \div 4 &= \frac{13}{4} \div 4 \\ &= \frac{13}{16}. \end{aligned}$$

Ex. 2. Divide £7. 8s. $1\frac{4}{11}$ d. by 6.

$$\begin{array}{r}
 \text{£.} \quad \text{s.} \quad \text{d.} \\
 6 \overline{) 7 \quad 8 \quad 1\frac{4}{11}} \\
 \underline{1 \quad 4 \quad 8\frac{8}{33}}
 \end{array}
 \qquad
 1\frac{5}{11} \div 6 = \frac{16}{11} \div 6$$

$$= \frac{16}{66} = \frac{8}{33}.$$

13. *To reduce a compound fraction to a simple one.*

DEF. A compound fraction is the fraction of a fraction.


Rule. Multiply the numerators together for a new numerator, and the denominators together for a new denominator.

Thus, $\frac{2}{3}$ of $\frac{4}{5} = \frac{8}{15}$.

For $\frac{1}{3}$ of $\frac{4}{5}$ is the same as $\frac{4}{5} \div 3$, and is therefore equal to $\frac{4}{15}$ by Art. 10. Therefore $\frac{2}{3}$ of $\frac{4}{5}$, which must be twice as great, will equal $\frac{8}{15}$.

Or we may prove it independently thus :

In order to obtain $\frac{2}{3}$ of $\frac{4}{5}$, we must divide each of our four fifths into 3 parts, and take two of them. But if we



divide a fifth into three parts, each resultant part (see figure) is a fifteenth. Taking then two of these parts we get $\frac{2}{15}$.

As then each fifth gives us $\frac{2}{15}$, $\frac{4}{5}$ will give us $\frac{8}{15}$. There-

fore $\frac{2}{3}$ of $\frac{4}{5} = \frac{8}{15}$.

$$\begin{aligned} \text{Ex. } 3\frac{5}{9} \text{ of } 2\frac{1}{4} \text{ of } \frac{3}{28} \text{ of } 4\frac{1}{5} &= \overset{2}{\cancel{3}\frac{5}{9}} \text{ of } \overset{2}{\cancel{2}\frac{1}{4}} \text{ of } \overset{3}{\cancel{3}\frac{3}{28}} \text{ of } \overset{3}{\cancel{4}\frac{1}{5}} \\ &= \frac{18}{5} \\ &= 3\frac{3}{5}. \quad \text{Ans.} \end{aligned}$$

"Cancelling," which we have adopted above, is nothing but another name for reducing a fraction to its lowest terms, and consequently the rule for cancelling is to divide numerator and denominator by the same numbers.

13'. To reduce fractions to their least common denominator.

We have proved that we may *divide* the numerator and denominator of a fraction by the same number, without altering its value; and it is clear that we may *multiply* the numerator and denominator of a fraction by the same number, without altering its value. Thus, $\frac{6}{9} = \frac{2}{3}$, and also $\frac{2}{3} = \frac{6}{9}$.

Suppose we wish to reduce the fraction $\frac{4}{7}$ to the denominator 42. Since the denominator 42 is derived from the denominator 7 by multiplying 7 by 6 ($42 \div 7 = 6$), the new numerator will be derived from the old numerator 4, by multiplying it by 6. Therefore $\frac{4}{7} = \frac{24}{42}$.

Hence we get the rule for reducing fractions to their least common denominator.

RULE. Find the L.C.M. of all the denominators (as that is the *smallest* number that *all* the denominators will divide exactly), and take this for the common denominator; for the new numerators, multiply each numerator by the quotient obtained by dividing the common denominator by its own denominator (as these quotients will be the numbers by which we have multiplied the respective denominators).

The object of this rule is to add together, or to subtract from each other, fractions of different denominators. We know that 3 apples + 5 apples = 8 apples; 3 pears + 5 pears = 8 pears; but we do not know how to express more simply 3 apples + 5 pears, because these are of different denominations. If, however, we were told that 3 apples were equal to 15 nuts, and 5 pears to 20 nuts, we should then know that 3 apples + 5 pears would be equal to 35 nuts, because we have now reduced the original denominations to the same denomination. So in fractions, we know that $\frac{3}{7} + \frac{5}{7}$ (3 sevenths + 5 sevenths) = $\frac{8}{7}$, and that $\frac{3}{9} + \frac{5}{9} = \frac{8}{9}$; but we do not at once know what $\frac{3}{7} + \frac{5}{9}$ are, because these are of different denominations. But by the above rule,

$$\frac{3}{7} = \frac{3 \times 9}{63} = \frac{27}{63}; \quad \frac{5}{9} = \frac{5 \times 7}{63} = \frac{35}{63};$$

therefore $\frac{3}{7} + \frac{5}{9} = \frac{27}{63} + \frac{35}{63} = \frac{62}{63}$, because we have now reduced the original denominations to the same denomination.

Similar remarks may evidently be applied to subtraction.

14. *Addition of fractions.*

Ex. Simplify $2\frac{3}{4}$ of $3\frac{2}{3} + \frac{111}{16} + 2\frac{4}{5}$ of $4\frac{1}{8}$ of $1\frac{3}{8} + 4\frac{2}{3}$ of $\frac{2}{15}$ of $2\frac{1}{8}$ of $1\frac{3}{7}$.

$$2\frac{3}{4} \text{ of } 3\frac{2}{3} = \frac{11}{4} \text{ of } \frac{11}{3} = \frac{121}{12} = 10\frac{1}{12}.$$

$$\frac{111}{16} = 5\frac{15}{16}.$$

$$2\frac{4}{5} \text{ of } 4\frac{1}{8} \text{ of } 1\frac{3}{8} = \frac{\cancel{14}^7}{5} \text{ of } \frac{\cancel{14}^7}{8} \text{ of } \frac{11}{8} = \frac{2541}{160} = 15\frac{141}{160}.$$

$$4\frac{2}{3} \text{ of } \frac{2}{15} \text{ of } 2\frac{1}{8} \text{ of } 1\frac{3}{7} = \frac{\cancel{14}^7}{3} \text{ of } \frac{\cancel{2}}{\cancel{15}^3} \text{ of } \frac{17}{8} \text{ of } \frac{\cancel{14}^7}{\cancel{7}^2} = \frac{17}{9} = 1\frac{8}{9}.$$

$$\begin{aligned} \therefore \text{Ans.} &= 10\frac{1}{12} + 5\frac{15}{16} + 15\frac{141}{160} + 1\frac{8}{9} \\ &= 31\frac{120 + 1350 + 1269 + 1280}{1440} \\ &= 31\frac{4019}{1440} \\ &= 33\frac{1139}{1440}. \end{aligned}$$

N.B. If it is preferred, the following method of doing the above example may be adopted.

$$\begin{aligned}
& 2\frac{3}{4} \text{ of } 3\frac{2}{3} + \frac{111}{16} + 2\frac{4}{5} \text{ of } 4\frac{1}{8} \text{ of } 1\frac{3}{8} + 4\frac{2}{3} \text{ of } \frac{2}{15} \text{ of } 2\frac{1}{8} \text{ of } 1\frac{3}{7} \\
&= \frac{11}{4} \text{ of } \frac{11}{3} + \frac{111}{16} + \frac{7}{5} \text{ of } \frac{33}{8} \text{ of } \frac{11}{8} + \frac{2}{3} \text{ of } \frac{2}{15} \text{ of } \frac{17}{8} \text{ of } \frac{10}{7} \\
&= \frac{121}{12} + \frac{111}{16} + \frac{2541}{160} + \frac{17}{9} \\
&= 10\frac{1}{12} + 5\frac{15}{16} + 15\frac{141}{160} + 1\frac{8}{9} \\
&= 31\frac{120 + 1350 + 1269 + 1280}{1440} \\
&= 31\frac{4019}{1440} \\
&= 33\frac{1139}{1440}.
\end{aligned}$$

The same method might be applied in Ex. 3, Art. 15, instead of the one there employed, and in other Arts.

15. *Subtraction of fractions.*

Always work with mixed numbers as in Addition. When this is not possible, proceed as follows :

$$\text{Ex. 1. } 7\frac{1}{4} - 2\frac{5}{6} = 5\frac{3-10}{12} = 4\frac{15-10}{12} = 4\frac{5}{12}. \quad \text{Ans.}$$

$$\text{Ex. 2. } 9 - 2\frac{3}{5} = 7 - \frac{3}{5} = 6\frac{5-3}{5} = 6\frac{2}{5}. \quad \text{Ans.}$$

In cases of many additions and subtractions in the same sum, *never insert brackets*. They are useless. The best rule is ; add the fractions or numbers that have to be

added, and add those that have to be subtracted; then subtract the latter result from the former.

Ex. 3. Simplify

$$5\frac{1}{5} - 2\frac{5}{6} - 3\frac{3}{10} + \frac{13}{2} - 16\frac{1}{4} + 3\frac{1}{12} + 8\frac{1}{8}.$$

$$\begin{aligned} 5\frac{1}{5} + \frac{13}{2} + 3\frac{1}{12} + 8\frac{1}{8} &= 5\frac{1}{5} + 6\frac{1}{2} + 3\frac{1}{12} + 8\frac{1}{8} \\ &= 22 \frac{24 + 60 + 10 + 15}{120} \end{aligned}$$

$$= 22 \frac{109}{120};$$

$$2\frac{5}{6} + 3\frac{3}{10} + 16\frac{1}{4} = 21 \frac{50 + 18 + 15}{60}$$

$$= 21 \frac{83}{60}$$

$$= 22 \frac{23}{60};$$

$$\therefore \text{Ans.} = 22 \frac{109}{120} - 22 \frac{23}{60} = \frac{109 - 46}{120}$$

$$= \frac{63}{120}$$

$$= \frac{21}{40}.$$

16. *Multiplication of fractions.* See Note A, and the next article.

17. *Division of fractions.* See Note B.

The whole theory of multiplication and division may

be dealt with and illustrated in many different manners, more or less simple. Some of these will be discussed in the notes at the end of the volume. Whatever plan of explanation be finally followed, it is clear, however, that for beginners, the theory should be made as short and simple as possible, and all complex, or even exact reasoning, on a subject so confessedly difficult, be left, till they have attained a much riper age. The following would seem to be sufficient at first.

DEF. *If we multiply the numerators of two fractions together for a new numerator, and the denominators together for a new denominator, we are said to multiply the fractions together.*

Since division reverses multiplication, the rule for division of fractions will be *to invert the divisor, and multiply.*

18. Since by the above rule of division of fractions

$$5 \div 7 = \frac{5}{1} \div \frac{7}{1} = \frac{5}{1} \times \frac{1}{7} = \frac{5}{7},$$

we get a new definition of a fraction, namely, *the numerator divided by the denominator.*

19. *To find the value of a fraction of a concrete quantity.*

Some latitude of working should be allowed here. However, a very general rule can be given *not to reduce to lowest denominations.* This is almost always a tedious and unnecessary method. We will give a few examples.

Ex. 1. Find the value of £7. 9s. $8\frac{1}{2}d. \times 3\frac{2}{5}$.

$$\begin{array}{r}
 \text{£.} \quad \text{s.} \quad \text{d.} \\
 7 \quad 9 \quad 8\frac{3}{4} \\
 \hline
 3 \\
 22 \quad 9 \quad 2\frac{1}{4} \\
 \hline
 2 \quad 19 \quad 10\frac{7}{10} \\
 \hline
 25 \quad 9 \quad 0\frac{12}{10} \quad \text{Ans.}
 \end{array}$$

$$\begin{array}{r}
 \text{£.} \quad \text{s.} \quad \text{d.} \\
 7 \quad 9 \quad 8\frac{3}{4} \\
 \hline
 2 \\
 5 \quad \overline{14 \quad 19 \quad 5\frac{1}{2}} \\
 2 \quad 19 \quad 10\frac{7}{10}
 \end{array}$$

Ex. 2. Find the value of £7. 9s. $8\frac{3}{4}d. \div 5\frac{3}{5}$.

$$\begin{array}{r}
 \text{£.} \quad \text{s.} \quad \text{d.} \\
 7 \quad 9 \quad 8\frac{3}{4} \div 5\frac{3}{5}, \text{ i.e. } \div \frac{28}{5}, \text{ i.e. } \times \frac{5}{28}. \\
 5 \\
 28 \quad \overline{37 \quad 8 \quad 7\frac{1}{8}} \\
 1 \quad 6 \quad 8\frac{1}{8} \quad \text{Ans.}
 \end{array}$$

$$\begin{array}{r}
 23 \frac{1}{3} \div 28 = \frac{70}{3} \times \frac{1}{28} \\
 \phantom{23 \frac{1}{3} \div 28 =} \frac{10}{3} \times \frac{1}{7} \\
 \phantom{23 \frac{1}{3} \div 28 =} = \frac{10}{21} \\
 \phantom{23 \frac{1}{3} \div 28 =} = \frac{5}{6}.
 \end{array}$$

Ex. 3. Find the value of £7. 9s. $8\frac{3}{4}d. \div \frac{7}{26}$.

$$\begin{array}{r}
 \text{£.} \quad \text{s.} \quad \text{d.} \\
 7 \quad 9 \quad 8\frac{3}{4} \div \frac{7}{26}, \text{ i.e. } \times \frac{26}{7}, \text{ i.e. } \times 3\frac{5}{7}. \\
 3 \\
 22 \quad 9 \quad 2\frac{1}{4} \\
 \hline
 5 \quad 6 \quad 11\frac{1}{8} \\
 \hline
 27 \quad 16 \quad 1\frac{9}{8} \quad \text{Ans.}
 \end{array}$$

$$\begin{array}{r}
 \text{£.} \quad \text{s.} \quad \text{d.} \\
 7 \quad 9 \quad 8\frac{3}{4} \\
 \hline
 5 \\
 7 \quad \overline{37 \quad 8 \quad 7\frac{1}{8}} \\
 5 \quad 6 \quad 11\frac{1}{8}
 \end{array}$$

Ex. 4. Find the value of $3\frac{7}{11}$ of $\frac{10}{7}$ of $\frac{77}{540}$ of 27s.

$$\begin{aligned}
 & \frac{3 \frac{7}{11}}{4 \frac{2}{7}} \text{ of } \frac{10 \frac{5}{7}}{7 \frac{1}{2}} \text{ of } \frac{77}{540} \text{ of } 27s. \\
 &= \frac{\cancel{40}^4}{\cancel{44}^{\cancel{11}}_3} \times \frac{\cancel{7}}{\cancel{30}^{\cancel{10}}_3} \text{ of } \frac{\cancel{15}^5}{\cancel{7}} \times \frac{2}{\cancel{15}} \text{ of } \frac{\cancel{77}^7}{\cancel{540}^{\cancel{60}}_9} \text{ of } \frac{\cancel{27}^9}{\cancel{1}} s. \\
 &= \frac{14}{3} s. \\
 &= 4 \frac{2}{3} s. \\
 &= 4s. 8d. \quad \text{Ans.}
 \end{aligned}$$

20. To reduce one quantity (or the fraction of one quantity) to the fraction of another (or of the fraction of another).

Ex. 1. Reduce 3 cwt. 2 qrs. 3 lbs. to the fraction of a ton.

cwt.	qrs.	lbs.	ton.
3	2	3	1
<hr style="width: 100%;"/>			<hr style="width: 100%;"/>
4			20
14 qrs.			20 cwt.
<hr style="width: 100%;"/>			<hr style="width: 100%;"/>
28			112
<hr style="width: 100%;"/>			<hr style="width: 100%;"/>
115			2240 lbs.
<hr style="width: 100%;"/>			
28			
<hr style="width: 100%;"/>			
395		lbs.	

$$\begin{aligned}
 \therefore 3 \text{ cwt. } 2 \text{ qrs. } 3 \text{ lbs.} &= \frac{395}{2240} \text{ of } 1 \text{ ton} \\
 &= \frac{79}{448} \text{ of } 1 \text{ ton.} \quad \text{Ans.}
 \end{aligned}$$

The reason of this is clear. For, since 1 lb. is $\frac{1}{2240}$ of 2240 lbs., therefore 395 lbs. is $\frac{395}{2240}$ of 2240 lbs., i.e. 3 cwt. 2 qrs. 3 lbs. is $\frac{395}{2240}$ (or $\frac{79}{448}$) of 1 ton.

Ex. 2. Reduce $\frac{3}{8}$ of $1\frac{1}{2}$ of 10s. $7\frac{1}{2}d.$ to the fraction of £4. 4s. $4\frac{1}{2}d.$

s. d.	£. s. d.
10 $7\frac{1}{2}$	4 4 $4\frac{1}{2}$
<u>12</u>	<u>20</u>
127d.	84s.
<u>2</u>	<u>12</u>
255 half-pence.	1012d.
	<u>2</u>
	2025 half-pence.

$\therefore \frac{3}{8}$ of $1\frac{1}{2}$ of 10s. $7\frac{1}{2}d.$ = $\frac{3}{8}$ of $1\frac{1}{2}$ of $\frac{255}{2025}$ of £4. 4s. $4\frac{1}{2}d.$

$$\begin{aligned}
 &= \frac{3}{8} \text{ of } \frac{3}{2} \text{ of } \frac{17}{240} \text{ of } £4. 4s. 4\frac{1}{2}d. \\
 &= \frac{17}{240} \text{ of } £4. 4s. 4\frac{1}{2}d. \quad \text{Ans.}
 \end{aligned}$$

Here again, since 10s. $7\frac{1}{2}d.$ = $\frac{255}{2025}$ of £4. 4s. $4\frac{1}{2}d.$,
(by the 1st Ex.), therefore $\frac{3}{8}$ of $1\frac{1}{2}$ of 10s. $7\frac{1}{2}d.$ = $\frac{3}{8}$ of $1\frac{1}{2}$

of $\frac{255}{2025}$ of £4 4s. 4½d., which, worked out as above, gives us the result $\frac{17}{240}$ of £4 4s. 4½d.

Ex. 3. Reduce $5\frac{4}{5}$ of £2. 2s. 6d. to the fraction of $5\frac{1}{3}$ of £1. 9s.

£. s. d.

2 2 6

20

42s.

2

85 sixpences.

£. s.

1 9

20

29s.

2

58 sixpences.

$$\therefore 5\frac{4}{5} \text{ of } £2. 2s. 6d. = 5\frac{4}{5} \text{ of } \frac{85}{5\frac{1}{3} \text{ of } 58} \text{ of } \left(5\frac{1}{3} \text{ of } £1. 9s.\right)$$

$$= \frac{29}{5} \text{ of } \frac{85}{58} \times \frac{3}{16} \text{ of } \left(5\frac{1}{3} \text{ of } £1. 9s.\right)$$

$$= \frac{51}{32} \text{ of } \left(5\frac{1}{3} \text{ of } £1. 9s.\right)$$

$$= 1\frac{19}{32} \text{ of } \left(5\frac{1}{3} \text{ of } £1. 9s.\right). \text{ Ans.}$$

Ex. 4. Reduce $\frac{7}{9}$ of £1 - $\frac{2}{5}$ of 21s. to the fraction of 10s. 6d.

$$\frac{7}{9} \text{ of } £1 - \frac{2}{5} \text{ of } 21s. = \frac{\frac{7}{9} \text{ of } 20 - \frac{2}{5} \text{ of } 21}{10\frac{1}{2}} \text{ of } 10s. 6d.$$

$$\begin{aligned}
 \text{Now } \frac{\frac{7}{9} \text{ of } 20 - \frac{2}{5} \text{ of } 21}{10 \frac{1}{2}} &= \frac{\frac{140}{9} - \frac{42}{5}}{10 \frac{1}{2}} \\
 &= \frac{15 \frac{5}{9} - 8 \frac{2}{5}}{10 \frac{1}{2}} \\
 &= \frac{7 \frac{25-18}{45}}{10 \frac{1}{2}} \\
 &= \frac{7 \frac{7}{45}}{10 \frac{1}{2}} \\
 &= \frac{\overset{46}{\cancel{322}}}{45} \times \frac{2}{\cancel{21} \times 3} \\
 &= \frac{92}{135};
 \end{aligned}$$

$$\therefore \frac{7}{9} \text{ of } £1 - \frac{2}{5} \text{ of } 21s. = \frac{92}{135} \text{ of } 10s. 6d. \quad \text{Ans.}$$

21. *Rule of precedence of the four simple rules in Arithmetical questions (with or without brackets).*

Multiplication and division precede addition and subtraction.

If any of the quantities are in brackets, of course each bracket (being a single quantity with respect to the rest

of the expression) must be simplified first, before the above rule applies to the whole quantity to be simplified.

Ex. 1. $1 + 2 \times 3 + 4 = 1 + 6 + 4 = 11.$

Ex. 2. $(1 + 2) \times 3 + 4 = 3 \times 3 + 4 = 9 + 4 = 13.$

Ex. 3. $1 + 2 \times (3 + 4) = 1 + 2 \times 7 = 1 + 14 = 15.$

Ex. 4. $(1 + 2) \times (3 + 4) = 3 \times 7 = 21.$

DECIMALS.

22. By means of the radix ten, we are able to express any integers from unity *upwards* without limit. By an extension of the method adopted in numeration and notation, we are able to get a series of "decimal fractions" from unity *downwards* without limit. The series of decimal fractions must have ten, or some power* of ten, for their denominators. The turning-points in this extension of the old system are as follows :

.....	
thousands	
hundreds.....	$1000 \div 10 = 100,$
tens.....	$100 \div 10 = 10,$
units	$10 \div 10 = 1,$
tenths	$1 \div 10 = \frac{1}{10},$
hundredths.....	$\frac{1}{10} \div 10 = \frac{1}{100},$
thousandths	$\frac{1}{100} \div 10 = \frac{1}{1000},$
.....	

* See Note to Art. 27.

Now in the old system for integers, in such a number as 1234, since that system commenced from a fixed point, unity, we know that the 4 of the above number represents units, the 3, tens, the 2, hundreds, and the 1, a thousand, but since the extension of the system for fractions does not begin at a fixed point, nor end at a fixed point, we must know the value of some particular digit, as unity, in order to know the real position and value of the rest. This is done by placing a point after the unit, commonly called a decimal point, as thus; 123·4567. Here then the 3 represents units, the 2, tens, the 1, a hundred; whereas, on the other hand, the 4 represents tenths, the 5, hundredths, the 6, thousandths, the 7, ten-thousandths (or tenths of thousandths). We may tabulate these results in the following manner.

A figure in the first place before the decimal point represents units.

A figure in the second place before the decimal point represents tens, ($\text{unity} \times 10$).

A figure in the third place before the decimal point represents hundreds, ($\text{ten} \times 10$).

A figure in the fourth place before the decimal point represents thousands, ($\text{hundred} \times 10$).

.....

Again :—

A figure in the first place after the decimal point represents tenths, ($\text{unity} \div 10$).

A figure in the second place after the decimal point represents hundredths, ($\text{tenth} \div 10$).

A figure in the third place after the decimal point represents thousandths, ($\text{hundredth} \div 10$).

A figure in the fourth place after the decimal point represents ten-thousandths, (thousandth $\div 10$).

.....

Thus the number 123·4567 would be read, one hundred and twenty three, decimal 4567; where "decimal 4567" would mean 4 tenths, 5 hundredths, 6 thousandths, and 7 ten-thousandths.

Again, the number ·007 would be read, decimal 007, and would mean 7 thousandths, i. e., $\frac{7}{1000}$.

Conversely, $\frac{7}{100} = \cdot 07$.

23. From the above article, the *rules for reducing a decimal to a fraction, and a fraction with some power of ten for its denominator to a decimal*, may be at once deduced.

$$\begin{aligned} 21\cdot0347 &= 20 + 1 + \frac{3}{100} + \frac{4}{1000} + \frac{7}{10000} = 21 + \frac{300 + 40 + 7}{10000} \\ &= 21 \frac{347}{10000}. \end{aligned}$$

Conversely,

$$\frac{10051}{10000} = 1 + \frac{50}{10000} + \frac{1}{10000} = 1 + \frac{5}{1000} + \frac{1}{10000} = 1\cdot0051.$$

Again, $11 \frac{5}{1000} = 11\cdot005$.

Hence we are in a position to give the verbal results.

To reduce a decimal to a fraction.

RULE. Write down the figures after the decimal point, for the numerator, and 1 followed by as many cyphers as there are decimal places, for the denominator.

To reduce a fraction, whose denominator is some power of ten, to a decimal fraction.

RULE. Write down the numerator of the fraction, and mark off as many decimal places as there are cyphers in the denominator.

24. Any decimal is multiplied by 10, 100, 1000, &c. by moving the decimal point one, two, three, &c. decimal places to the right.

Any decimal is divided by 10, 100, 1000, &c. by moving the decimal point one, two, three, &c. decimal places to the left.

$$\text{Thus, } .012 \times 100 = \frac{12}{1000} \times \frac{100}{1} = \frac{12}{10} = 1.2.$$

$$\begin{aligned} \text{And } 1.234 \div 100 &= 1 \frac{234}{1000} \div 100 = \frac{1234}{1000} \times \frac{1}{100} \\ &= \frac{1234}{100000} = .01234. \end{aligned}$$

25. Multiplication of Decimals.

Leave out all superfluous decimal cyphers, thus:—

Ex. Multiply together .0003, 12.5, 50000, 20, .0004.

$$\begin{array}{r} .0003 \\ 12.5 \\ \hline .00375 \\ 50000 \\ \hline 187.50000 \\ 20 \\ \hline 3750.0 \\ .0004 \\ \hline 1.5000 \end{array}$$

Ans. = 1.5.

26. *Division of Decimals.*

It is as well that there should be but one rule learnt for the division of decimals. Perhaps the following will be found to be the clearest and the easiest to apply.

If the given divisor is not a whole number, make it so by removing its decimal point altogether, and shift the decimal point of the dividend as many places to the right as there were decimal figures in the divisor; annexing for this purpose, if necessary, decimal cyphers to the dividend.

Then, divide as if the given decimals were common integers; and when, in the process of division, the decimal point of the dividend is arrived at, place a decimal point in the quotient.

Decimal cyphers may be annexed to the dividend to any extent that may be required for carrying on the division.

Ex. Divide $\cdot 04$ by 20 , $\cdot 4$ by $\cdot 002$, $\cdot 0004$ by $\cdot 0002$, $\cdot 004$ by $\cdot 2$, 400 by $\cdot 02$.

27. *To reduce a fraction to a decimal.*

RULE. Divide the numerator by the denominator.

This follows from the 2nd definition of a vulgar fraction. Art. 18.

If the denominator of the fraction reduced to its lowest terms, consists only of powers* of two and five as factors,

* By the "power" of a number is meant the product of the number into itself any number of times. Thus 10 is said to be the first power

the decimal will (i) terminate. If it has no factors of two or five, the decimal will be (ii) a pure circulator. If it has factors of two or five, or both, *with other factors*, the decimal will be (iii) a mixed circulator.

For (i), leaving out of consideration the decimal point, a fraction is reduced to a decimal by multiplying the numerator by any power of 10 (i.e. by any powers of 2 and 5), and dividing the result by the denominator. Now, with each power of ten in the numerator, a factor of 2 and 5 will disappear (by cancelling) in the denominator. Also, when the less power of either 2 or 5 is exhausted, there will remain a power of 5 or 2. Each factor of this power will disappear by means of a further power of ten in the numerator. Hence, if the denominator of a fraction reduced to its lowest terms consists only of powers of two and five as factors, it is clear that the decimal will terminate.

It is also clear that the number of decimal places in the decimal will equal the number of factors of 2 or 5 in the denominator, whichever happens to be the greater. For every power of 10 introduced as a factor of the numerator gives one decimal place.

Again (ii), if the denominator has neither 2 nor 5 as a factor, it is clear that the decimal *will not terminate*. For none of the factors of the denominator can either disappear with those of the numerator (for the fraction is in its lowest terms) or with the powers of 2 and 5 (i.e. of 10) by which the numerator is multiplied. We have now to prove that the decimal will *circulate*.

In the division, we always bring down the same figure, namely, a cypher; hence, if ever at any step of the division of 10, 100 the second power of 10 (because it equals 10×10), 1000 the third power of 10 ($10 \times 10 \times 10$), &c.

an old remainder should be repeated, from that step the whole process will be repeated. But, since a remainder must be less than the divisor, the number of different remainders is limited. Therefore either the decimal will recur or will terminate. But we have proved that it does not terminate. Therefore it recurs.

Also it is clear from the above, that the number of decimal places in the period *at most* is one less than the denominator.

Lastly (iii), if the denominator of the fraction reduced to its lowest terms has factors of 2 or 5, or both, as well as other factors, the decimal will be a mixed circulator, the number of places before the period being equal to the power of 2 or 5 in the denominator, whichever happens to be greater, and the number of figures in the period being *at most* one less than the denominator when deprived of its factors of 2 and 5.

The truth of the last statement will be best seen from a few instances.

$$\frac{5}{6} \left(= \frac{5}{2 \times 3} \right) = \frac{50}{6} \div 10 = \frac{25}{3} \div 10 = 8\dot{3} \div 10 = \cdot 8\dot{3}.$$

$$\frac{5}{44} \left(= \frac{5}{4 \times 11} \right) = \frac{500}{44} \div 100 = \frac{125}{11} \div 100 = 11\dot{3}\dot{6} \div 100 = \cdot 11\dot{3}\dot{6}.$$

$$\begin{aligned} \frac{371}{1125} \left(= \frac{371}{125 \times 9} \right) &= \frac{371000}{1125} \div 1000 = \frac{2968}{9} \div 1000 \\ &= 329\dot{7} \div 1000 = \cdot 329\dot{7}. \end{aligned}$$

$$\frac{4}{35} \left(= \frac{4}{5 \times 7} \right) = \frac{40}{35} \div 10 = \frac{8}{7} \div 10 = 1\dot{1}4285\dot{7} \div 10 = \cdot 1\dot{1}4285\dot{7}.$$

The object of the form given to the fractions in the brackets is to indicate the reason of selection of the ensuing powers of ten, by which we multiply numerator and denominator.

The last example gives us the period as large as it can be, viz. one less than the denominator when deprived of its factors of 2 and 5.

All the above results, with others of somewhat less importance, and greater difficulty of arithmetical proof, are tabulated in Art. 44.

28. *Proof of the rule for reducing a circulating decimal to a fraction.*

First, for a pure circulator, as $\dot{3}\dot{7}$.

$$\begin{aligned}\dot{3}\dot{7} \times 100 &= 37\dot{3}\dot{7} \\ &= 37 + \dot{3}\dot{7};\end{aligned}$$

$$\therefore \dot{3}\dot{7} \times 99 = 37;$$

$$\therefore \dot{3}\dot{7} = \frac{37}{99}.$$

This agrees with the rule.

Next, for a mixed circulator, as $3\dot{4}\dot{7}$.

$$\begin{aligned}3\dot{4}\dot{7} \times 10 &= 3\dot{4}\dot{7} \\ &= 3\frac{47}{99} \text{ (by the above)} \\ &= \frac{3 \times 99 + 47}{99} \\ &= \frac{3(100 - 1) + 47}{99} \\ &= \frac{347 - 3}{99};\end{aligned}$$

$$\therefore 3\dot{4}\dot{7} = \frac{347 - 3}{99}.$$

This also agrees with the rule.

$$29. \quad \dot{9} = \frac{9}{9} = 1, \quad \cdot\dot{0}9 = \frac{9}{90} = \frac{1}{10} = \cdot 1, \quad \cdot 00\dot{9} = \frac{9}{900} \\ = \frac{1}{100} = \cdot 01, \text{ \&c.}$$

Hence, generally, the period 9 at any place of decimals may be replaced by a *one* of the preceding place. Thus we can assert at once that

$$\cdot 4\dot{9} = \cdot 5, \quad \cdot 74\dot{9} = \cdot 75, \quad \cdot 124\dot{9} = \cdot 125, \text{ \&c.}$$

30. It is well to remember that

$$1 \text{ farthing} = \frac{1}{4} = \cdot 25.$$

$$1 \text{ halfpenny} = \frac{1}{2} (= \cdot 50) = \cdot 5.$$

$$3 \text{ farthings} = \frac{3}{4} = \cdot 75.$$

$$\text{And of course, conversely, } \cdot 25 = \frac{1}{4}, \quad \cdot 75 = \frac{3}{4}, \quad \cdot 5 = \frac{1}{2}.$$

31. *To find the value of a decimal of any given quantity.*

It is well, in finding the value of a decimal of a given quantity, when the decimal is a circulator, *if the period is a long one, generally* to reduce to a fraction, if short, to work in decimals. We will give an example of the latter.

Ex. Find the value of $2\cdot 14\dot{5}$ of $5s. 8\frac{1}{2}d.$

$$\begin{array}{r} 5s. 8\frac{1}{2}d. \\ \frac{12}{68d.} \\ \frac{4}{275 \text{ farthings.}} \end{array}$$

$$\begin{array}{r}
 2'14545 \\
 \underline{275} \\
 1072727 \\
 1501818,1 \\
 \underline{429090,90} \\
 4 \overline{) 589'99999} \\
 12 \overline{) 147'49999} \\
 12 - 3
 \end{array}$$

$$\text{Ans.} = 12s. 3'49d. = 12s. 3'5d. \text{ (by Art. 29)} = 12s. 3\frac{1}{2}d.*$$

On examining the above, it will be observed that the proper corrections are introduced throughout, arising from the recurrence of the period. This of course is necessary to ensure a right result. Observe also that the *whole result*, 589'999..... farthings is divided by 4, as we never give answers in decimals, or fractions, *of farthings*.

N.B. The answer above in farthings is 589'9, i.e. 590 farthings = 12s. 3 $\frac{1}{2}$ d. This is a rather shorter and better method than the one adopted. As however we wished to shew a *general* method, and this immediate simplification only applied to the particular case, we purposely avoided it. No other period but 9 could have been thus simplified.

32. It will be found convenient in working decimals to remember the following result :

$$\frac{1}{7} = .14285\dot{7}.$$

Now the order of largeness of the above figures is 1, 2, 4, 5, 7, 8, and it will be found that if we begin with these figures respectively, and take the period in the same order, we shall get the values of the fractions

* The commas are introduced to point out where the process of multiplication *usually* stops.

$\frac{2}{7}, \frac{3}{7}, \frac{4}{7}, \frac{5}{7}, \frac{6}{7}$ respectively. Thus,

$$\frac{2}{7} = \cdot\dot{2}8571\dot{4},$$

$$\frac{3}{7} = \cdot\dot{4}2857\dot{1},$$

$$\frac{4}{7} = \cdot\dot{5}7142\dot{8},$$

$$\frac{5}{7} = \cdot\dot{7}1428\dot{5},$$

$$\text{and } \frac{6}{7} = \cdot\dot{8}5714\dot{2}.$$

Conversely, any one of these decimals may at once be reduced back again to a fraction by considering the order of merit of the figures of its period. Thus, $\cdot\dot{5}7142\dot{8} = \frac{4}{7}$, because 5 is the *fourth largest figure* of the six figures of the period. This is not arbitrary, but follows from Arts. 27 and 44, since the period contains the greatest number of figures possible, viz. one less than the denominator.

Example :

$$\cdot\dot{2}8571\dot{4} \text{ of } £30 + 6 \cdot\dot{8}5714\dot{2}£ + \cdot\dot{6} \text{ of } \cdot\dot{7}1428\dot{5} \\ \text{of } \cdot\dot{6}£ + 1\cdot\dot{3} \text{ of } \cdot\dot{4}2857\dot{1}s.$$

$$= \frac{2}{7} \text{ of } £30 + 6 \frac{6}{7}£ + \frac{2}{3} \text{ of } \frac{5}{7} \text{ of } \frac{3}{5}£ + 1 \frac{1}{3} \text{ of } \frac{3}{7}s.$$

$$= 8 \frac{4}{7}£ + 6 \frac{6}{7}£ + \frac{2}{7}£ + \frac{4}{3} \text{ of } \frac{3}{7}s.$$

$$= 15 \frac{5}{7}£ + \frac{4}{7}s.$$

$$= £15. 14 \frac{2}{7} s. + \frac{4}{7} s.$$

$$= £15. 14 \frac{6}{7} s.$$

$$= £15. 14s. 10 \frac{2}{7} d. \quad \text{Ans.}$$

32'. Reduce to a decimal accurate to 5 places,

$$16 \left(\frac{1}{5} - \frac{1}{3 \cdot 5^3} + \frac{1}{5 \cdot 5^5} - \frac{1}{7 \cdot 5^7} + \dots \right) - \frac{4}{239} *.$$

$$\frac{1}{5} = \cdot 2,$$

$$\frac{1}{5^3} = \cdot 04,$$

$$\frac{1}{5^5} = \cdot 008;$$

$$\therefore \frac{1}{3 \cdot 5^3} = \cdot 00266666.$$

$$\frac{1}{5^7} = \cdot 0016,$$

$$\frac{1}{5^9} = \cdot 00032,$$

$$\frac{1}{5^{11}} = \cdot 000064,$$

$$\frac{1}{5^{13}} = \cdot 0000128;$$

$$\therefore \frac{1}{7 \cdot 5^7} = \cdot 00000182.$$

$$\frac{1}{5^{15}} = \cdot 00000256,$$

$$\frac{1}{5^{17}} = \cdot 000000512;$$

$$\therefore \frac{1}{9 \cdot 5^9} = \cdot 00000005.$$

.....

.....

$$\frac{1}{5} = \cdot 2,$$

* The *dot* in the denominators implies multiplication. 5^3 denotes $5 \times 5 \times 5$; 5^5 , $5 \times 5 \times 5 \times 5 \times 5$; and so on; so that 5^3 is the third power, 5^5 , the fifth power, of 5, and so on.

$$\frac{1}{5 \cdot 5^8} = \cdot 000064,$$

$$\frac{1}{3 \cdot 5^8} = \cdot 00266666.$$

$$\frac{1}{9 \cdot 5^9} = \cdot 00000005,$$

$$\frac{1}{7 \cdot 5^7} = \cdot 00000182.$$

$$\therefore \frac{1}{5} + \frac{1}{5 \cdot 5^8} + \frac{1}{9 \cdot 5^9} = \cdot 20006405; \quad \therefore \frac{1}{3 \cdot 5^8} + \frac{1}{7 \cdot 5^7} = \cdot 00266848.$$

$$\therefore \frac{1}{5} - \frac{1}{3 \cdot 5^8} + \frac{1}{5 \cdot 5^8} - \frac{1}{7 \cdot 5^7} + \dots = \cdot 19739557;$$

$$\therefore 16 \left(\frac{1}{5} - \frac{1}{3 \cdot 5^8} + \frac{1}{5 \cdot 5^8} - \frac{1}{7 \cdot 5^7} + \dots \right) = 3 \cdot 15832912.$$

$$\frac{4}{239} = \cdot 01673640;$$

$$\therefore 16 \left(\frac{1}{5} - \frac{1}{3 \cdot 5^8} + \frac{1}{5 \cdot 5^8} - \frac{1}{7 \cdot 5^7} + \dots \right) - \frac{4}{239} = 3 \cdot 14159272;$$

$$\therefore \text{Ans.} = 3 \cdot 14159.$$

PROPORTION.

33. Proportion is usually taught by the method of "statement." There are several advantages arising from the use of this method. It is the neatest, it is the shortest, it can be as easily taught as any other method *with respect to getting out right answers*, it is the method universally required from a business man, so that it would be a disgrace not to know how to "state a rule of three sum." There is also the "common sense" method which has this inestimable advantage, that those who use it understand the reason of the steps, which we cannot, we are afraid, generally predicate of those who use the statement method.

As, moreover, Arithmetic should appeal to the reason, and not merely to the memory, it would seem well to *teach* the common sense method, and afterwards indicate the method by "statement," by going through a few examples. We will work a sum by the rule of common sense.

Ex. If 30 cwt. are carried 15 miles for £5. 8s. 9d., how far ought 80 cwt. to be carried for £29?

If 30 cwt. for 1305d. are carried 15 miles;

∴ 1 cwt. for 1305d. is carried 15 ÷ 30 miles;

∴ 1 cwt. for 1d. is carried $\frac{15 \times 30}{1305}$ miles;

∴ 80 cwt. for 1d. are carried $\frac{15 \times 30}{1305 \times 80}$ miles;

∴ 80 cwt. for 6960d. are carried $\frac{15 \times 30 \times 6960}{1305 \times 80}$ miles;

$$\therefore \text{Ans.} = \frac{\overset{30}{\overset{3}{15} \times \overset{3}{30} \times \overset{6960}{6960}}}{\underset{267}{\underset{87}{\underset{29}{1305}} \times 80}} \text{ miles} = 30 \text{ miles.}$$

The rules to be observed are :

1. Write down what you are told.
2. Write it down in such a manner that that comes last which is the same kind of thing as the answer.

In the statement method, the rules are :

1. The third term must be the same kind of thing as the answer.
2. The other two terms (which are compared) must be of a like sort.

3. If the answer is greater than the third term, put the greater of the two like things in the second place; if less than the third term, put the less of the two like things in the second place.

4. Multiply the second and third terms together, and divide by the first. This will give the answer.

E.g. If 5 yds. of cloth cost 12s., what would 15 yds. cost?

$$\begin{array}{cccc} \text{yds.} & \text{yds.} & \text{s.} & \text{s.} \\ 5 & : 15 & :: 12 & : \text{Ans.} \end{array}$$

$$\therefore \text{Ans} = \frac{15 \times 12}{5} \text{ s.} = 36\text{s.} = \text{£}1. 16\text{s.}$$

This is the right statement. The following, though it will bring out a right answer, has no meaning, and should never be allowed.

$$\begin{array}{cccc} \text{yds.} & \text{s.} & \text{yds.} & \text{s.} \\ 5 & : 12 & :: 15 & : \text{Ans.} \end{array}$$

It has no meaning, because no arithmetical comparison can be instituted between 5 yds. and 12s., &c.

We will give an example of double rule of three by the statement method. *No other statement should ever be allowed.*

If 12 persons spend £160 in 4 months, how many will £853. 6s. 8d. last for 8 months?

$$\begin{array}{ccccccc} \text{£.} & & \text{£.} & & & & \\ 160 & : & 853\frac{1}{3} & & & & \\ & & & & \text{per.} & \text{per.} & \\ & & & & :: 12 & : \text{Ans.} & \\ \text{mo.} & & \text{mo.} & & & & \\ 8 & : & 4 & & & & \end{array}$$

$$\begin{aligned}
 \therefore \text{Ans.} &= \frac{12 \times 853 \frac{1}{3} \times 4}{160 \times 8} \text{ persons} = \frac{\overset{2}{\cancel{12}} \times \overset{16}{\cancel{2560}} \times \cancel{4}}{\cancel{160} \times \underset{2}{\cancel{8}} \times \underset{2}{\cancel{3}}} \text{ persons} \\
 &= 32 \text{ persons.}
 \end{aligned}$$

RULES REDUCIBLE TO PROPORTION.

34. All the money rules, called Interest, Discount, Brokerage, Stocks, and Profit and Loss, are only proportion applied to particular subjects, which involve new definitions. Hence, when the definitions and nomenclature are thoroughly understood, these rules can contain no fresh difficulty.

35. *Simple Interest.*

This rule introduces the following words: Interest, Principal and Amount. The Rate of Interest is the interest on £100 for 1 year. Let us now solve two or three questions. See Note C.

Ex. 1. Find the amount on £250. 12s. 6d. from March 26, 1840, to Oct. 31, 1842, at 3 per cent.

From March 6, 1840, to Oct. 31, 1842, is 2 years, 219 days = $2 \frac{219}{365}$ years = $2 \frac{3}{5}$ years.

$$£250. 12s. 6d. = £250 \frac{5}{8} = £\frac{2005}{8}.$$

First, find the interest.

If £100 in 1 year gains £3;

$$\therefore \text{£1 in 1 year gains } \frac{\text{£}3}{100};$$

$$\therefore \text{£}\frac{2005}{8} \text{ in 1 year gains } \frac{\text{£}3 \times 2005}{100 \times 8};$$

$$\begin{aligned} \therefore \text{£}\frac{2005}{8} \text{ in } \frac{13}{5} \text{ years gains } & \frac{\text{£}3 \times 2005 \times 13}{100 \times 8 \times 5} \\ & = \text{£}\frac{15639}{800} = \text{£}19. 10s. 11\frac{7}{10}d. \text{ Int.} \end{aligned}$$

Add £250 12 6 Principal.

Hence we get £270. 3s. 5 $\frac{7}{10}$ d. Amt. Ans.

N.B. It will be found convenient, generally, if not always, to work in £'s and fractions of £'s as above, and not to reduce to lower denominations.

It is as well to observe that an easier plan of finding simple interest or amount can generally be adopted than the above, viz. by the ordinary rule, which is, to multiply the principal, the no. of years, and the rate per cent. together, and divide by 100. This will give us the interest. The reason of this rule will not require explanation. Thus, in the above example,

$$\begin{aligned} \text{Int.} = \text{£} \frac{250 \frac{5}{8} \times 3 \times 2 \frac{3}{5}}{100} &= \text{£} \frac{2005 \times 3 \times 13}{100 \times 8 \times 5}, \\ &= \dots\dots\dots \end{aligned}$$

The reason that we have not employed this method is partly because it is of limited application, and partly that it may be thoroughly seen *in all cases* that Interest and the other money rules are not new rules, but only various applications of proportion adapted to new definitions. See Art. 34.

It might be asked, "Why cannot we find the amount at once in the last sum, beginning thus :

If £100 in 1 year amounts to £103, &c.?"

The answer is, that no proportion is true which involves *at the same time* years and amount. This can easily be proved, for it is clear that, though interest increases in the same proportion as the length of years, amount does not, because a part of it (*viz.* the principal) remains stationary during the time. Thus, £100 at 5 per cent. in 1 year amounts to £105; in 2 years to £110, *not to* £210.

Ex. 2. At what rate will £220. 12s. 6d. become £240. 4s. 8 $\frac{2}{3}$ d. in $3\frac{1}{3}$ years?

Here, £240. 4s. 8 $\frac{2}{3}$ d. is amt. But years and amount cannot exist together in a proportion sum. Subtract therefore the principal from the amount to get the interest. Thus, we have £220. 12s. 6d. gains £19. 12s. 2 $\frac{2}{3}$ d. in $3\frac{1}{3}$ years.

$$£220. 12s. 6d. = £220 \frac{5}{8} = £1765 \frac{5}{8}.$$

$$£19. 12s. 2\frac{2}{3}d. = £19 \frac{11}{18} = £353 \frac{11}{18}.$$

$$\text{If } £1765 \frac{5}{8} \text{ in } \frac{10}{3} \text{ years gains } £353 \frac{11}{18};$$

$$\therefore £1 \text{ in } \frac{10}{3} \text{ years gains } £ \frac{353 \times 8}{18 \times 1765};$$

$$\therefore £1 \text{ in 1 year gains } £ \frac{353 \times 8 \times 3}{18 \times 1765 \times 10};$$

The question is thus reduced to the following: If £100 amounts to £124 $\frac{1}{2}$, what sum will amount to £49. 9s. 5½d.? This is a *proportion* sum, as we have got rid of *years*.

$$£49. 9s. 5\frac{1}{2}d. = £49 \frac{7}{320} = £\frac{15687}{320}.$$

If £124 $\frac{1}{2}$ is the amount on £100;

$$\therefore £1 \text{ is the amount on } £\frac{100 \times 2}{249};$$

$$\therefore £\frac{15687}{320} \text{ is the amount on } £\frac{100 \times 2 \times 15687}{249 \times 320}$$

$$\begin{array}{r} 63 \\ 5229 \\ 15687 \\ \times 249 \\ \hline 12687 \\ 24900 \\ 156870 \\ \hline 382233 \end{array}$$

$$= £\frac{315}{8} = £39\frac{3}{8}$$

$$= £39. 7s. 6d. \text{ Ans.}$$

N.B. The practical rule for solving interest questions involving amount is thus found to be: If possible, get rid of amount, by means of the data (i.e. by subtracting interest or principal from amount). If not possible, find the amount on £100 for the given time at the given rate.

36. Compound Interest.

Here the process adopted by us all through the latter rules may be employed. But a far shorter method is to take the ordinary rule and work by decimals.

Ex. 1. Find the amount of £95. 16s. 8d. for 2 years, at $2\frac{1}{2}$ per cent., comp. int.

£.	s.	d.
95	16	8
		$2\frac{1}{2}$
191	13	4
47	18	4
239	11	8
20		
791		
12		
1100		

∴ Amt. for 1st year = £95. 16s. 8d. + £2. 7s. 11d.
= £98. 4s. 7d.

£.	s.	d.
98	4	7
		$2\frac{1}{2}$
196	9	2
49	2	$3\frac{1}{2}$
245	11	$5\frac{1}{2}$
20		
911		
12		
137 $\frac{1}{2}$		

$$37\frac{1}{2} \div 100 = \frac{37\frac{1}{2}}{2} \times \frac{1}{100} = \frac{3}{8}.$$

∴ Amt. for 2 years = £98. 4s. 7d. + £2. 9s. 1 $\frac{3}{4}$ d.
= £100. 13s. 8 $\frac{1}{2}$ d. Ans.

Ex. 2. Find the comp. int. on £225 for 3 years,
at $3\frac{3}{4}$ per cent.

$$\begin{array}{r} £225 \\ \quad 3\frac{3}{4} \\ \hline 675 \\ 168\cdot75 \\ \hline 843\cdot75 \end{array}$$

\therefore Amt. for 1st year = £225 + £8·4375 = £233·4375.

$$\begin{array}{r} £233\cdot4375 \\ \quad 3\frac{3}{4} \\ \hline 7003125 \\ 175078125 \\ \hline 875\cdot390625 \end{array}$$

\therefore Amt. for 2 years = £233·4375 + £8·75390625
= £242·19140625.

$$\begin{array}{r} £242\cdot19140625 \\ \quad 3\frac{3}{4} \\ \hline 72657421875 \\ 1816435546875 \\ \hline 908\cdot2177734375 \end{array}$$

\therefore Amt. for 3 years = £242·19140625 + £9·082177734375
= £251·273583984375.

\therefore Ans. = £251·273583984375 - £225 = £26·273583984375
= £26. 5s. 5 $\frac{169}{256}$ d.

This method might have been adopted in Ex. 1. We should have to begin thus :

$$£95. 16s. 8d. = £95\frac{5}{6} = £95\cdot8\dot{3} = £95\cdot83333\ldots$$

Corrections on account of the recurrence would have to be introduced at the different steps.

37. *Discount.*

This rule introduces the new words, discount, bill, present value.

Suppose a man borrows £500, with promise to pay £550 in 3 years time. The £550 is called the bill, and is the amount that is due to the money-lender at the end of 3 years. The £500 is called (very naturally) the present value, and the difference (*viz.* £50), the discount.

Supposing we want to find the present value of £550, due at the end of 3 years at 5 per cent., the question resolves itself into the old interest question, what principal in 3 years at 5 per cent. will become £550. For the money-lender might be supposed to say to the borrower, "It makes no difference to me whether you pay me £550 in 3 years (as we agreed), or whether you pay me *now* such a sum of money that if I put it into the bank at 5 per cent., I shall find at the end of 3 years that it has become £550." Hence a "bill" is only another name for amount, present value for principal, and discount for interest.

Ex. Find the discount on a bill of £419. 12s. 1d., drawn March 6, at 7 months, discounted Sept. 15, at 5 per cent.

Here the bill was drawn March 6, at 7 months, and was therefore due Oct. 9 (allowing for the three days of grace). Now it was discounted Sept. 15, i.e. 24 days before it was due. Hence we have the question :

What principal in 24 days will become £419. 12s. 1d., at 5 per cent. ?

$$£419. 12s. 1d. = £419 \frac{29}{48} = £\frac{20141}{48}.$$

$$\begin{aligned} \text{Int. on } £100 \text{ at } 5 \text{ per cent. for } 24 \text{ days} &= £\frac{24}{365} \text{ of } \frac{£}{1} \\ &= £\frac{24}{73}; \end{aligned}$$

$$\therefore \text{Amt.} = £100 \frac{24}{73} = £\frac{7324}{73}.$$

If $£\frac{7324}{73}$ is amount on £100;

$$\therefore £1 \text{ is amount on } £\frac{100 \times 73}{7324};$$

$$\therefore £\frac{20141}{48} \text{ is amount on } £\frac{100 \times 73 \times 20141}{7324 \times 48}$$

$$\begin{aligned} &= £\frac{20075}{48} = £418 \frac{11}{48} \\ &= £418. 4s. 7d. \end{aligned}$$

$$\therefore \text{Discount} = £419. 12s. 1d. - £418. 4s. 7d. = £1. 7s. 6d. \text{ Ans.}$$

38. *Stocks.*

This rule introduces the words, capital, stock, income, and such new expressions as, "the 5 per cents. are at 90." This expression means that £5 is the interest on £100 stock, which is worth £90.

All questions in stocks proper may be reduced to three or four primitive questions. We will solve a few; in each case supposing that the 5 per cents. are at 90.

Ex. 1. What is the value of £1000 stock?

If £100 stock is worth £90;

$$\therefore \text{£1 stock is worth } \text{£} \frac{90}{100};$$

$$\begin{aligned} \therefore \text{£1000 stock is worth } \text{£} \frac{90 \times 1000}{100} \\ = \text{£}900. \text{ Ans.} \end{aligned}$$

Ex. 2. How much stock can be obtained for £1000?

If £90 will buy £100 stock;

$$\therefore \text{£1 will buy } \text{£} \frac{100}{90} \text{ stock};$$

$$\begin{aligned} \therefore \text{£1000 will buy } \text{£} \frac{100 \times 1000}{90} \text{ stock} \\ = \text{£}1111 \frac{1}{9} \text{ stock. Ans.} \end{aligned}$$

Ex. 3. What is my income from £1000 stock?

If £100 stock produces £5;

$$\therefore \text{£1 stock produces } \text{£} \frac{5}{100};$$

$$\begin{aligned} \therefore \text{£1000 stock produces } \text{£} \frac{5 \times 1000}{100} \\ = \text{£}50. \text{ Ans.} \end{aligned}$$

Ex. 4. What is my income from investing £1000?

If £90 produces £5;

$$\therefore \text{£1 produces } \text{£} \frac{5}{90};$$

$$\begin{aligned} \therefore \text{£1000 produces } \text{£} \frac{5 \times 1000}{90} \\ = \text{£}55 \frac{5}{9} \\ = \text{£}55. 11s. 1\frac{1}{3}d. \text{ Ans.} \end{aligned}$$

This last is done in the shortest, but not the most natural way. Properly speaking, we ought first to have found the amount of stock purchased, as in Ex. 2. This is $\text{£}1111\frac{1}{9}$ stock. Then proceed in the same manner as in Ex. 3, thus:

If $\text{£}100$ stock produces $\text{£}5$;

$\therefore \text{£}1$ stock produces $\text{£}\frac{5}{100}$;

$\therefore \text{£}\frac{10000}{9}$ stock produces $\text{£}\frac{5 \times 10000}{100 \times 9}$
 $= \text{£}55. 11s. 1\frac{1}{3}d.$ Ans.

The best method of doing examples in stocks is to separate each question into the events that, as a matter of fact, really happened, and consider these events consecutively, e.g.

Ex. 5. A person invests $\text{£}18150$ in the 3 per cents. at $90\frac{3}{4}$, and, on their rising to 91, transfers to the $3\frac{1}{2}$ per cents. at $97\frac{1}{2}$; what increase does he make thereby in his annual income?

First Question. He buys stock with $\text{£}18150$; find the amount of stock bought.

If $\text{£}90\frac{3}{4}$ buys $\text{£}100$ stock;

$\therefore \text{£}1$ buys $\text{£}\frac{100 \times 4}{363}$ stock;

$$\begin{array}{r}
 50 \\
 550 \\
 6050 \\
 6600 \\
 7150 \\
 \hline
 363 \\
 777 \\
 777 \\
 \hline
 = £20000 \text{ stock.}
 \end{array}$$

$\therefore £18150 \text{ buys } £ \frac{100 \times 4 \times 18150}{363} \text{ stock}$

Second Question. This £20000 stock he sells, when the stock is at 91. Find how much money he gets for his stock.

If £100 stock is sold for £91;

$$\therefore £1 \text{ stock is sold for } £ \frac{91}{100};$$

$$\begin{array}{r}
 \therefore £20000 \text{ stock is sold for } £ \frac{91 \times 20000}{100} \\
 \hline
 = £18200.
 \end{array}$$

Third Question. With this £18200, he buys some more stock, at $97\frac{1}{2}$. Find the amount of stock bought.

If $£97\frac{1}{2}$ buys £100 stock;

$$\therefore £1 \text{ buys } £ \frac{100 \times 2}{195} \text{ stock};$$

$$\begin{array}{r}
 280 \\
 3540 \\
 4050 \\
 4600 \\
 5150 \\
 \hline
 39 \\
 3 \\
 \hline
 = £ \frac{56000}{3} \text{ stock} \\
 = £18666\frac{2}{3} \text{ stock.}
 \end{array}$$

$\therefore £18200 \text{ buys } £ \frac{100 \times 2 \times 18200}{39} \text{ stock}$

Since we are asked to find the difference in his income caused by these transactions, we must find his *first* income, when he had £20000 stock in the three per cents., and his *second* income, when he had £18666 $\frac{2}{3}$ stock in the $3\frac{1}{2}$ per cents. Hence,

Fourth Question. If £100 stock produces £3;

$$\therefore \text{£1 stock produces } \text{£} \frac{3}{100};$$

$$\begin{aligned} \therefore \text{£20000 stock produces } \text{£} \frac{3 \times 20000}{100} \\ = \text{£600.} \end{aligned}$$

Fifth Question. If £100 stock produces £ $\frac{7}{2}$;

$$\therefore \text{£1 stock produces } \text{£} \frac{7}{2 \times 100};$$

$$\begin{aligned} \therefore \text{£} \frac{56000}{3} \text{ stock produces } \text{£} \frac{7 \times 56000}{2 \times 100 \times 3} \\ = \text{£} \frac{1960}{3} \\ = \text{£653. 6s. 8d.} \end{aligned}$$

Sixth Question. Hence the difference in income
= £653. 6s. 8d. - £600

= £53. 6s. 8d. Ans.

Sometimes, when no ambiguity can arise, the word stock is not expressed. Thus, if it is said that a man bought £1000, sold £1000, transferred £1000 from the 4 per cents. at 90 to the 5 per cents. at 80, or had £1000 in the 5 per cents. at 90, in all these cases the £1000 is clearly stock. If it is said that a man laid out £1000, or invested £1000 in

the 4 per cents. at 90, or that a sum in the 4 per cents. at 90 was worth £1000; in all these cases the £1000 is clearly capital.

39. Profit and Loss.

The words here used are cost price, selling price, gain (or loss). It must be remembered that a man always *gains or loses on the cost price*. Thus a loss of 5 per cent. means a loss of £5 on a thing *costing* £100, so that the selling price would be £95.

Ex. 1. If tea be bought at 6s. 3d. per lb., and sold at 5s. 7d., what is the loss per cent.?

Here loss on the cost price 6s. 3d. is 8d.

If on 75d. the loss is 8d.

$$\therefore \text{on } 1d. \text{ the loss is } \frac{8}{75}d.;$$

$$\therefore \text{on } 100d. \text{ the loss is } \frac{8 \times 100}{75}d.$$

$$= \frac{32}{3}d. = 10\frac{2}{3}d.;$$

$$\therefore \text{Ans. is } 10\frac{2}{3} \text{ per cent.}$$

Obs. Since we are comparing 6s. 3d. and 100, they must be of the same kind (pence suppose), and since by gain per cent. we mean so much on 100 of the same denomination, it follows that, our answer being $10\frac{2}{3}$ pence

on 100 pence, we get at once the result $10\frac{2}{3}$ per cent., whatever be the denomination.

Ex. 2. If $5\frac{1}{2}$ per cent. be gained by selling butter at £5. 5s. 6d. per cwt., what will be the gain per cent., by selling it at 1s. 3d. per lb.?

As we gain and lose on the *cost price*, it will be necessary, from the first part of the question, to find the cost price, in order to solve the latter part of the same. First then, we will find the cost price.

$$£5. 5s. 6d. = £5\frac{11}{40} = £\frac{211}{40}.$$

If a thing which is sold for $£105\frac{1}{2}$ costs £100;

∴ a thing which is sold for £1 costs $£\frac{100 \times 2}{211}$;

∴ a thing which is sold for $£\frac{211}{40}$ costs $£\frac{5 \times 100 \times 2 \times 211}{211 \times 40}$
= £5.

If 1 lb. is sold for 1s. 3d.;

∴ 1 cwt. is sold for $\frac{5}{4} \times \frac{28}{1}$ s. = 140s. = £7;

∴ gain on the cwt. is £7 - £5 = £2.

If on £5 the gain is £2;

∴ on £1 the gain is $\frac{£2}{5}$;

∴ on £100 the gain is $£\frac{2 \times 100}{5}$
= £40;

∴ Ans. = 40 per cent.

The above may be done a shorter way, but it is unsatisfactory, as the ordinary student would probably find it difficult to explain the reason of his method. It is well to solve all questions in Profit and Loss by the single simple principle that all gain or loss is on the cost price. No method but the above should be allowed. All other methods are dangerous.

Ex. 3. If a person purchase pins when they are 18 in a row, and sell them 11 in a row at the same price, how much is his gain per cent.?

As a person gains and loses on the *cost price*, we must imagine a cost price, to work the question simply and clearly.

Suppose 18 pins to cost 18 pence;

\therefore 11 pins are sold for 18 pence.

But since 18 pins cost 18 pence;

\therefore 11 pins cost 11 pence;

\therefore 11 pins cost 11 pence and are sold for 18 pence;

\therefore 7d. is the gain on the cost price 11d.

If on 11d. the gain is 7d.;

\therefore on 1d. the gain is $\frac{7}{11}d.$

\therefore on 100d. the gain is $\frac{7 \times 100}{11}d.$

$= 63\frac{7}{11}d.$

\therefore Ans. $= 63\frac{7}{11}$ per cent.

40. *Proportional parts.*

This rule really only includes an easy class of questions involving knowledge of fractions and proportion, and there is no reason why it should usurp a high place in Arithmetic.

If a sum of money is to be divided among A and B in the ratio of 3 : 5, it is clear that as often as A has £3, B will have £5. Therefore, out of every £8 that is divided between them, A has £3 and B has £5. Therefore A will have $\frac{3}{8}$ of the whole, B $\frac{5}{8}$ of the whole.

We will solve two questions.

Ex. 1. A and B engaged in trade, their capitals being in the ratio of 4 : 5 ; and, at the end of 3 months, they withdrew respectively $\frac{2}{3}$ and $\frac{3}{4}$ of their capitals : how should they divide their whole gain, £335, at the end of the year ?

Suppose A to have £4 ;

$\therefore B$ will have £5.

Since A withdraws $\frac{2}{3}$ of his capital, he will have $\frac{1}{3}$ of £4 ($= \frac{£4}{3}$) remaining.

Since B withdraws $\frac{3}{4}$ of his capital, he will have $\frac{1}{4}$ of £5 ($= \frac{£5}{4}$) remaining.

Now a capital of £4 for 3 months is equivalent to £12 for 1 month.

And a capital of $\pounds \frac{4}{3}$ for 9 months is equivalent to $\pounds 12$ for 1 month;

\therefore (adding) we may represent *A*'s capital at $\pounds 24$ for 1 month.

Again, a capital of $\pounds 5$ for 3 months is equivalent to $\pounds 15$ for 1 month.

And a capital of $\pounds \frac{5}{4}$ for 9 months is equivalent to $\pounds 11 \frac{1}{4}$ for 1 month;

\therefore (adding) we may represent *B*'s capital at $\pounds 26 \frac{1}{4}$ for 1 month.

Hence, if we divide the profits, $\pounds 335$, in the ratio of the two capitals, $\pounds 24$ and $\pounds 26 \frac{1}{4}$, we shall get the proper answer.

(Observe, a capital of $\pounds 4$ for 3 months is equivalent to a capital of $\pounds 12$ for 1 month, for by an *equivalent* capital we mean a capital that will produce the same profits. Of course if $\pounds 4$ were the *profits* in 3 months, the profits in 1 month would be $\pounds \frac{4}{3}$.)

Ex. 2. If 8 oz. of gold, 10 carats fine, and 2 oz., 11 carats fine, be mixed with 6 oz. of unknown fineness, and that of the mixture be 12 carats, what was the unknown fineness?

By 8 oz. of gold, 10 carats fine, we mean that $\frac{10}{24}$ of the 8 oz. is gold, the rest being alloy, see Art. 4. In this and all similar questions, reduce to, and work in ordinary Tr. weight, and if the answer is to be given in carats, re-

duce to 24ths of the whole mass in question. The number of 24ths will give us the fineness required.

8 oz. of gold, 10 carats fine, contain $\frac{80}{24}$ oz. of pure gold.

2 oz. of gold, 11 carats fine, contain $\frac{22}{24}$ oz. of pure gold.

\therefore (adding) 10 oz. of the mixed gold contain $\frac{102}{24}$ oz. of pure gold.

But the whole 16 oz. of the mixed gold, 12 carats fine, contain $\frac{192}{24}$ oz. of pure gold;

\therefore the 6 oz. of gold introduced contain $\frac{90}{24}$ oz. of pure gold;

\therefore 1 oz. of gold introduced contains $\frac{15}{24}$ oz. of pure gold;

\therefore the unknown fineness is 15 carats. Ans.

(For in every oz. of the new gold we have proved that 15 parts out of 24 are pure gold.)

41. If A can do $\frac{3}{5}$ of a piece of work in 7 days, $\frac{1}{2}$ of which A and B together can do in 5, how long would B alone be doing $\frac{5}{7}$ of it?

Since A can do $\frac{3}{5}$ of the work in 7 days;

\therefore he can do the whole work in $\frac{5}{3}$ of $7 = \frac{35}{3}$ days;

\therefore he can do $\frac{3}{35}$ of the work in 1 day.

Since A and B can do $\frac{1}{2}$ of the work in 5 days;

\therefore they can do the whole work in 10 days;

\therefore they can do $\frac{1}{10}$ of the work in 1 day;

$\therefore B$ alone can do $\frac{1}{10} - \frac{3}{35}$ of the work in 1 day;

that is, $\frac{1}{70}$ of the work in 1 day.

If B can do $\frac{1}{70}$ of the work in 1 day;

\therefore he can do the whole work in 70 days;

\therefore he can do $\frac{5}{7}$ of the work in $\frac{5}{7}$ of 70 days;

that is, in 50 days. Ans.

42. If two fruit women were willing to exchange 12 apples for 49 pears, 121 nuts for 25 pears, 6 oranges for 5 plums, 154 nuts for 45 oranges; how many apples would they be willing to exchange for 77 plums?

12 apples = 49 pears,

25 pears = 121 nuts,

154 nuts = 45 oranges,

6 oranges = 5 plums,

77 plums = x apples.

$$\therefore \text{Ans.} = \frac{4 \quad 5 \quad 14 \quad 2 \quad 11}{\cancel{12} \times \cancel{25} \times \cancel{154} \times \cancel{6} \times \cancel{77}} \text{ apples}$$

$$\frac{49 \times \cancel{121} \times \cancel{45} \times 5}{\cancel{7} \quad \cancel{11} \quad 9 \quad 3}$$

$$= 16 \text{ apples. Ans.}$$

We shall not explain the reason of the above. The student may easily verify it by a series of proportions. It is called the "chain rule," and is useful in working questions of "Exchange" in a ready and simple manner. It will be observed in the two upright columns *that each of them contains all the different denominations.*

43. *Per centage.*

A large number of the latter examples, after proportion, may be worked out in a different manner by a careful consideration of the words "per centage."

Thus, 2 per cent. means that 2 articles out of 100 of the same articles are taken into consideration; that is, we are considering $\frac{2}{100}$ of the whole.

If a man loses 5 per cent. of his money, in any way, it means that he loses $\frac{5}{100}$ of his money. He will have remaining $\frac{95}{100}$. We can thus get rid of proportion, in examples of per centage, and reduce per centage to fractions.

On the whole, however, we object to this method, because ambiguities are likely to arise from the looseness with which the expression "so much per cent." is used in common language. To make questions of this sort free from ambiguity, we should either have to define scientifically popular language, or explain carefully the exact sense in which we intend to use the words in the different cases.

44. We will finish with certain practical rules which will be found *very useful* for checking errors, and in many

ways stimulating the ease and rapidity of correct work. We do not propose to prove them generally, or explain on what principles they depend. Some may be proved by simple Arithmetic, others by Arithmetic diluted with Algebra; while the proof of others belongs entirely to the region of Algebra. But the results are important and easy to be remembered.

a. A number is divisible by 2, 4, 8, respectively, if the number represented by its last 1, 2, 3, figures respectively is divisible by 2, 4, 8, respectively.

The rule for 4 can be proved thus :

100 is divisible by 4, therefore any number of hundreds is divisible by 4; if then a number is divisible by 4, the remainder found by removing the hundreds, that is, the number represented by the last two digits, must be divisible by 4. So for 8, 16, 32 ...

A number is divisible by 5, if it ends in a 5 or a cypher.

A number is divisible by 10, if it ends in a cypher.

A number is divisible by 3, 9, respectively, if the sum of its digits is divisible by 3, 9, respectively. See Note D.

A number is divisible by 11, if the difference of the sums of its digits in the odd and even places is divisible by 11. See Note D.

The rule for 7 is more complex than ordinary division by 7.

The rule for 6 is the combination of the rules for 2 and 3; for 12, the combination of the rules for 3 and 4; &c. Care must be taken to split 12 into its factors 4, 3 (not 6, 2 for instance); 24 into its factors 8, 3; 75 into its factors 25, 3, &c.

N.B. These rules are useful, in reducing fractions to their lowest terms, in cancelling, L.C.M., G.C.M., &c.

β . The shortest way of multiplying a number by 25 is, to add two cyphers and divide by 4.

The shortest way of multiplying a number by 125 is, to add three cyphers and divide by 8.

So, the shortest way of dividing a number by 25 is, to mark off two decimal places, and multiply by 4.

And the shortest way of dividing a number by 125 is, to mark off three decimal places, and multiply by 8.

N.B. It may be observed that an odd number is divisible by 25 if it ends in 25 or 75; by 125 if it ends in 125, 625, 375, 875.

These rules are useful in reducing fractions to decimals, and conversely; generally in cases where the decimals terminate. They are also useful in other work.

γ . Let us take a fraction *in its lowest terms*, which we wish to reduce to a decimal. Then the following rules hold.

1. If the denominator of the fraction consists *only* of powers of 2 and 5 as factors, the resulting decimal will terminate. Art. 27.

2. If the denominator of the fraction has no factor of 2 or 5, the resulting decimal will be a pure circulator. Art. 27.

3. If the denominator of the fraction consists partly of powers of 2 or 5, or both, as factors, and partly of other factors, the resulting decimal will be a mixed circulator. Art. 27.

4. The number of figures in the terminating decimal, in (1), will be that power of 2 or 5 in the denominator of the fraction which happens to be the greater. Art. 27.

5. The number of figures in the period of the decimal, in (2), will be a measure of some measure diminished by unity of the denominator of the fraction.

6. The number of figures *before* the period of the decimal, in (3), will follow rule (4); *in* the period of the decimal, will follow rule (5). Art. 27.

7. When the decimal is a pure circulator, and the period a full period, the same period will occur, only commencing at a different figure of the period, for every fraction with the same denominator, but a different numerator. See Art. 32.

8. When the decimal is a pure circulator, and the period a full period, every figure in the last half of the period will be a complement to 9 of every corresponding figure in the first half of the period.

[N.B. By a full period, we mean a period that is the largest possible, i.e., one less than the denominator of the fraction. See Rule (5)].

We will exemplify rules (7) and (8). We shall find that

$$\frac{1}{19} = \cdot\dot{0}5263157894736842\dot{1}.$$

Here (2) and (5) are satisfied, and the number of figures in the period being $19 - 1$, it is a full period. Hence we can find $\frac{2}{19}$, $\frac{3}{19}$, &c., at once. Thus:

$$\frac{2}{19} = \cdot\dot{1}05263157894736842\dot{2},$$

$$\frac{3}{19} = \cdot\dot{1}57894736842105263\dot{3},$$

$$\frac{15}{19} = \cdot\dot{7}8947368421052631\dot{5}.$$

We began this last $\cdot\dot{7}89\dots$ because there were only three larger fractions than $\frac{15}{19}$, viz. $\frac{16}{19}$, $\frac{17}{19}$, $\frac{18}{19}$; and there were only 3 decimals beginning with higher figures than $\cdot\dot{7}89\dots$, viz. $\cdot\dot{8}42\dots$, $\cdot\dot{8}94\dots$ and $\cdot\dot{9}47\dots$. So for any other numerator. In Art. 32, we have already discussed the important fractions $\frac{1}{7}$, $\frac{2}{7}$, &c., in like manner.

Again, take one of the above fractions, as $\frac{15}{19}$, and divide the period by a mark into its two halves, thus:

$$\frac{15}{19} = \cdot\dot{7}89473684,21052631\dot{5}.$$

Hence (8) is clearly seen to be true, for $7 + 2 = 9$, $8 + 1 = 9$, $9 + 0 = 9$, $4 + 5 = 9$, &c. So for any other numerator, or for any of the fractions $\frac{1}{7}$, $\frac{2}{7}$, &c., in Art. 32.

Ex. The first 29 figures of the period, for the fraction $\frac{1}{59}$, are found to be

$$\cdot\dot{0}1694915254237288135593220338\dots\dots$$

Complete the period.

There are other very curious properties about the remainders, in reducing fractions to decimals, and about the periods, both when full and incomplete, but we do not give them here, partly because they are not so important as the above, partly because it is not easy to express them sufficiently generally or concisely for practical work.

8. 6s. 8d. is $\frac{1}{3}$ of a £; 2s. 6d. is $\frac{1}{8}$ of a £.

If we wish to reduce shillings, pence, and farthings to the fraction of a £, the denominator of the resulting fraction *can have no other factors* than 3, 5, and powers of 2.

NOTES.

NOTE A. *Multiplication of fractions.*

By "multiplication" we mean the repetition of a number a certain number of times. Thus, 7×4 means 7 repeated 4 times, or $7 + 7 + 7 + 7$, or 28. So 7 apples multiplied by 4 is 28 apples. From this definition it is clear that the multiplier must be abstract and a whole number. According to our definition then, it is clear that we could not multiply by any of the following quantities, 4 apples, $\frac{4}{5}$, $+4$, -4 . Of course, on the other hand, we

could multiply any of them, 4 apples, $\frac{4}{5}$, $+4$, -4 , by any abstract integer. It is no matter what the multiplicand may be, it is only necessary that the multiplier should be of a certain character.

If then we are to talk for the future of multiplying by such things as 4 apples, $\frac{4}{5}$, $+4$, -4 , it can only be by altering the original meaning of the word multiplication, or at least extending and generalising it. This is done repeatedly in algebra, and thus we get the well-known results,

$$+a \times +b = +ab,$$

$$+a \times -b = -ab,$$

$$-a \times +b = -ab,$$

$$-a \times -b = +ab.$$

And again,

$$a^m \times a^n = a^{m+n}, \quad a^m \div a^n = a^{m-n}, \quad (a^m)^n = a^{mn},$$

whether m and n be positive, negative, fractional, integral, or zero.

If again, in Arithmetic, we allow ourselves to use the expression, 4 ft. \times 5 ft. = 20 sq. ft., (as in Art. 5) it can either be by supposing that expression to be the abbreviation for the proposition "if we multiply the number of feet in the length of any rectangle by the number of feet in its breadth, we shall get the number of square feet in its area," as is fully explained in the same Article, or we must suppose the meaning of multiplication to be defined anew to suit the case, something for instance as follows.

"A square foot is said to be obtained geometrically by multiplying the foot of its length by the foot of its breadth." Hence it will be seen (as in Art. 5) that the area of any rectangle would be obtained by multiplying its length by its breadth.

Again, in fractions, we do use such expressions as $\frac{4}{7} \times \frac{5}{9}$, $4 \times \frac{5}{7}$, &c. This is called multiplication of fractions. Of course then here also we must redefine multiplication. (N.B. It is clear it will not be necessary to do so for such cases as $\frac{5}{7} \times 4$.)

The new definition may be given in various ways. We will here state and discuss a definition.

DEF. We are said to multiply two fractions together if we multiply the numerators for a new numerator, and the denominators for a new denominator.

This seems at first sight a very arbitrary definition. Any one has a right to ask, Whence did such a definition arise?

To shew this, let us take a case, e.g. $\frac{3}{7} \times \frac{4}{5}$.

$\frac{3}{7} \times 4 = \frac{12}{7}$. But we have to multiply not by 4, but by $\frac{4}{5}$, i.e. by $\frac{1}{5}$ of 4. Hence *it will be convenient to agree* that $\frac{3}{7} \times \frac{4}{5}$, or $\frac{3}{7} \times \frac{1}{5}$ of 4 shall mean $\frac{1}{5}$ of $\left(\frac{3}{7} \times 4\right)$, i.e. $\frac{1}{5}$ of $\frac{12}{7}$; which can be easily proved to be $\frac{12}{35}$ (Art. 13). This gives us the ordinary Arithmetical rule.

The important thing here is not to mistake the above for a proof. *It is no proof at all.* It is a convenient assumption, or convention, or new definition of the word "multiplication." Any other definition might have been taken, but this suits us best. The italicised words must be remembered: "it will be convenient to agree."

It might still be asked, however, why the above assumption is more convenient than any other. This we will proceed to explain.

It is clear that $\frac{3}{7} \times 4 = \frac{12}{7}$. Again, it is clear that $\frac{3}{7} \times \frac{12}{3} = \frac{36}{21}$. For $\frac{12}{3} = 4$, and therefore the case in point is $\frac{3}{7} \times 4$, which equals $\frac{12}{7}$, which equals $\frac{36}{21}$. Since also in the first instance, instead of 4 we may write $\frac{4}{1}$, it is clear that we shall not get a wrong answer in the case of

$\frac{3}{7} \times 4$, if for 4 we write either $\frac{4}{1}$ or $\frac{12}{3}$, and then multiply the numerators for a new numerator, and the denominators for a new denominator. In other words, this rule will apply whenever our multiplier is an integer, even if in the form of a fraction. This shews us why our definition is a convenient one when our multiplier is not an integer (in which case multiplication has lost its original meaning), because we shall thus have one rule for all cases; in other words, the form of the answer will not have to be altered to suit the different cases.

For instance, we thus know that $\frac{a}{b} \times \frac{c}{d} = \frac{ac}{bd}$ and have not to enquire what $\frac{c}{d}$ is. Otherwise we should have to say "If $\frac{c}{d}$ is really an integer, $\frac{a}{b} \times \frac{c}{d} = \frac{ac}{bd}$. If, however, $\frac{c}{d}$ is not an integer, $\frac{a}{b} \times \frac{c}{d} = \dots$ " whatever our definition of multiplication of fractions makes it equal.

As it is very important that the student should thoroughly understand that there is no meaning in the words "Prove the rule for the multiplication of fractions," or "Shew that $\frac{3}{7} \times \frac{4}{5} = \frac{12}{35}$," but only in "Illustrate the rule," or "Shew why we say that $\frac{3}{7} \times \frac{4}{5} = \frac{12}{35}$," or "Shew that $\frac{3}{7} \times \frac{4}{5}$ is not necessarily equal to $\frac{12}{35}$;" and as the quasi proof given above is often mistaken for a real proof, we will discuss another instance in the same sort of manner, and come to a conclusion different from that which is usually adopted.

5^7 means the repetition of 5 as a factor 7 times ;

$\therefore 5^{-7}$ means the repetition of 5 as a factor - 7 times ;

$\therefore 5^{-7}$ means the repetition of - 5 as a factor 7 times ;

$\therefore 5^{-7} = (-5)^7 = -78125.$

Now this result is not true. Not true, that is to say, *necessarily*. However unwise, it would not be wrong to give a definition of multiplication in the above case, which should make the above result true. But even if the new definition should give that result, the *proof* would still be false—indeed it is manifest there would be no proof possible ; only the application of the new definition.

One more word then with regard to results.

$\frac{3}{7} \times 4$ is necessarily $= \frac{12}{7}.$

$\frac{3}{7} \times \frac{4}{5}$ is equal to $\frac{12}{35}$ by a convention governed by common sense, but perfectly arbitrary.

NOTE B. *Division of fractions.*

We now come to division of fractions. The rule given is—Invert the divisor and multiply. The question is—Can this rule be *proved* or no, or if it can, in what manner?

If we allow ourselves the two definitions of division given in Art. 3, this rule can be *proved*; thereby differing from multiplication. The proof will be as follows.

It is clear that $\frac{3}{5} \div 7 = \frac{3}{35}$ by the 2nd def. of division.

This agrees with the rule, for

$$\frac{3}{5} \div 7 = \frac{3}{5} \div \frac{7}{1} = \frac{3}{5} \times \frac{1}{7} = \frac{3}{35}.$$

Again, it is clear that $\frac{3}{5} \div \frac{7}{5} = \frac{3}{7}$. This is clear from the 1st definition of division. This also agrees with the rule, for

$$\frac{3}{5} \div \frac{7}{5} = \frac{3}{5} \times \frac{5}{7} = \frac{3}{7}.$$

Lastly, let us consider $\frac{3}{5} \div \frac{4}{7}$. This is the same as $\frac{21}{35} \div \frac{20}{35}$. The result of this is $\frac{21}{20}$ by the 1st definition of division. And by the rule also we obtain

$$\frac{3}{5} \div \frac{4}{7} = \frac{3}{5} \times \frac{7}{4} = \frac{21}{20}.$$

Hence, in all cases the rule is proved.

We do not like this proof, partly because objections might be raised against it as not compatible with the only definition of fractions we have at present thought it convenient to give, and more especially because we think it a better plan to treat division of fractions in the same manner as we have already treated multiplication of fractions. We leave it to the ingenuity of the student to do this.

Or we might illustrate the rule in the following manner.

Division reverses multiplication. Since then $\frac{3}{5} \times \frac{4}{7} = \frac{12}{35}$, it follows that $\frac{12}{35} \div \frac{4}{7} = \frac{3}{5}$. It is clear that this quotient may be obtained by inverting the divisor and multiplying. Hence the rule.

N.B. The whole theory of multiplication and division of fractions may be illustrated and dealt with in a different and interesting manner, by extending our definition of a fraction, and introducing axioms, or by making a separation between pure arithmetic and algebraical arithmetic, and laying down the results in parallel columns. It will however probably be felt that we have already dwelt on this subject quite sufficiently.

NOTE C. Every sum worked out in the text, from Art. 35 inclusive to the end, has been handled by the "common sense" method of proportion. We propose here to indicate the course of proof of the same sums, proceeding by the method of statement instead. We conceive that a student might reasonably be allowed to take his choice of either method to adopt.

Art. 35. Ex. 1. Find the amount on £250. 12s. 6d. from March 26, 1840, to Oct. 31, 1842, at 3 per cent.

From March 6, 1840, to Oct. 31, 1842, is 2 years, 219 days = $2 \frac{219}{365}$ years = $2 \frac{3}{5}$ years = $\frac{13}{5}$ years.

$$£250. 12s. 6d. = £250 \frac{5}{8} = £\frac{2005}{8}.$$

First, find the interest.

$$\begin{array}{rcl} \text{£.} & & \text{£.} \\ 100 & : & \frac{2005}{8} \\ \text{year.} & \text{years.} & :: 3 : \text{Int.} \\ 1 & : & \frac{13}{5} \end{array}$$

Hence the interest, and hence the amount.

Ex. 2. At what rate will £220. 12s. 6d. become £240. 4s. 8 $\frac{2}{3}$ d. in $3\frac{1}{3}$ years?

$$£220. 12s. 6d. = £220\frac{5}{8} = £\frac{1765}{8}.$$

$$\begin{aligned}\text{Int.} &= £240. 4s. 8\frac{2}{3}d. - £220. 12s. 6d. = £19. 12s. 2\frac{2}{3}d. \\ &= £19\frac{11}{18} = £\frac{353}{18}.\end{aligned}$$

$$\begin{array}{rcl}\begin{array}{c}£. \\ 1765 \\ 8\end{array} & : & \begin{array}{c}£. \\ 100\end{array} \\ \text{years. year.} & & \\ 3\frac{1}{3} : 1 & & \end{array}$$

$$\begin{array}{rcl}£. & & \\ 353 & : & \text{Ans.} \\ 18 & & \end{array}$$

∴ Required rate =

Ex. 3. What sum will become £49. 0s. 5 $\frac{1}{4}$ d. in $5\frac{1}{4}$ years, at $4\frac{2}{3}$ per cent.?

Interest on £100 for $5\frac{1}{4}$ years, at $4\frac{2}{3}$ per cent., will be

$$\begin{aligned}& £ \frac{100 \times 5\frac{1}{4} \times 4\frac{2}{3}}{100} \\ &= £ \frac{\overset{7}{\cancel{2}}\overset{7}{\cancel{2}}}{\underset{2}{4}} \times \frac{\overset{7}{\cancel{4}}\overset{7}{\cancel{4}}}{\underset{3}{3}} \\ &= £ \frac{49}{2} \\ &= £24\frac{1}{2};\end{aligned}$$

$$\therefore \text{Amount} = £124 \frac{1}{2} = £\frac{249}{2}.$$

$$£49. 9s. 5\frac{1}{2}d. = £49 \frac{7}{320} = £\frac{15687}{320};$$

$$\therefore \frac{\overset{£.}{249}}{2} : \frac{\overset{£.}{15687}}{320} :: 100 : \text{Ans.}$$

Hence the required principal.

Art. 36. Exs. 1 and 2. These do not require to be considered here, as they have not been done by proportion in the text.

Art. 37. Find the discount on a bill of £419. 12s. 1d., drawn March 6, at 7 months, discounted Sept. 15, at 5 per cent.

Here the bill was drawn March 6, at 7 months, and was therefore due Oct. 9 (allowing for the three days of grace). Now it was discounted Sept. 15, i.e. 24 days before it was due.

$$£419. 12s. 1d. = £419 \frac{29}{48} = £\frac{20141}{48}.$$

$$\begin{aligned} \text{Interest on } £100, \text{ at 5 per cent. for 24 days} &= £\frac{24}{365} \times \frac{5}{100} \\ &= £\frac{24}{73}; \end{aligned}$$

$$\therefore \text{Amt.} = £100 \frac{24}{73} = £\frac{7324}{73}.$$

$$\frac{\overset{£.}{7324}}{73} : \frac{\overset{£.}{20141}}{48} :: 100 : \text{Present value.}$$

Hence the present value, and hence the required discount.

It will be observed that it is not sufficient in the first two terms, to compare money merely with money, nor in the third term, to have it merely the same denomination as the answer. Care must be taken that the first two terms should be *both amounts*, and that the third term should be *present value*, the same as the answer.

N.B. If a bill fall due on a day that has no existence, as the 31st of a month of 30 days, or the 29th, 30th, or 31st of Feb. (in leap year, the 30th or 31st of Feb. only), it is considered to be *nominally* due on the last day of the month in question, and therefore *legally* due on the 3rd of the following month.

Art. 38. Ex. 1. What is the value of £1000 stock, in the 5 per cents. at 90?

$$\begin{array}{cccc} \text{£. st.} & \text{£. st.} & \text{£.} & \text{£.} \\ 100 & : & 1000 & :: 90 : \text{Ans.} \end{array}$$

Ex. 2. How much stock can be obtained for £1000, in the 5 per cents. at 90?

$$\begin{array}{cccc} \text{£.} & \text{£.} & \text{£. st.} & \text{£. st.} \\ 90 & : & 1000 & :: 100 : \text{Ans.} \end{array}$$

Ex. 3. What is my income from £1000 stock, in the 5 per cents. at 90?

$$\begin{array}{cccc} \text{£. st.} & \text{£. st.} & \text{£.} & \text{£.} \\ 100 & : & 1000 & :: 5 : \text{Ans.} \end{array}$$

Ex. 4. What is my income from investing £1000 in the 5 per cents. at 90?

$$\begin{array}{cccc} \text{£.} & \text{£.} & \text{£.} & \text{£.} \\ 90 & : & 1000 & :: 5 : \text{Ans.} \end{array}$$

N.B. Here the first two terms are *both capital*; the last two, *income*.

Ex. 5. A person invests £18150 in the 3 per cents. at $90\frac{3}{4}$, and, on their rising to 91, transfers to the $3\frac{1}{2}$ per cents. at $97\frac{1}{2}$; what increase does he make thereby in his annual income?

£. £. £. st.

$$90\frac{3}{4} : 18150 :: 100 : \text{stock purchased};$$

$$\therefore \text{Stock purchased} = £ \frac{\overset{50}{\cancel{18150}} \times 100 \times 4}{\begin{array}{r} 363 \\ 726 \\ 72 \end{array}} \text{ stock}$$

$$= £20000 \text{ stock.}$$

£. st. £. st. £.
100 : 20000 : 91 : cash received;

$$\therefore \text{Cash received} = £ \frac{91 \times 20000}{100}$$

$$= £18200.$$

£. £. £. st.

$$97\frac{1}{2} : 18200 :: 100 : \text{fresh stock purchased};$$

$$\therefore \text{fresh stock purchased} = £ \frac{\overset{280}{\cancel{100}} \times \overset{3640}{\cancel{18200}} \times 2}{\begin{array}{r} 195 \\ 39 \\ 3 \end{array}} \text{ stock}$$

$$= £ \frac{56000}{3} \text{ stock.}$$

$$\begin{array}{ccc} \text{£. st.} & \text{£. st.} & \text{£.} \\ 100 & : 20000 & :: 3 : \text{first income;} \end{array}$$

$$\begin{aligned} \therefore \text{first income} &= \text{£} \frac{3 \times 20000}{100} \\ &= \text{£}600. \end{aligned}$$

$$\begin{array}{ccc} \text{£. st.} & \text{£. st.} & \text{£.} \\ 100 & : \frac{56000}{3} & :: 3\frac{1}{2} : \text{second income;} \end{array}$$

$$\begin{aligned} \therefore \text{second income} &= \text{£} \frac{280 \times 56000 \times 7}{2 \times 3 \times 100} \\ &= \text{£} \frac{1960}{3} \\ &= \text{£}653. 6s. 8d.; \end{aligned}$$

$$\begin{aligned} \therefore \text{required difference of income} &= \text{£}653. 6s. 8d. - \text{£}600, \\ &= \text{£}53. 6s. 8d. \quad \text{Ans.} \end{aligned}$$

A single proportion might have been made to replace the third and fifth of the above proportions, as thus :

$$\begin{array}{ccc} \text{£.} & \text{£.} & \text{£.} \\ 97\frac{1}{2} & : 18200 & :: 3\frac{1}{2} : \text{second income.} \end{array}$$

We prefer the method adopted notwithstanding, as it exhibits the natural progression of the events, and puts the whole working in a clearer and more logical light.

Art. 39. Ex. 1. If tea be bought at 6s. 3d. per lb., and sold at 5s. 7d., what is the loss per cent. ?

Here loss on the cost price 6s. 3d. is 8d.

$$\begin{array}{ccc} d. & d. & d. \\ 75 & : 100 & :: 8 : \text{Ans.} \end{array}$$

N.B. Here the first two are cost price, the last two are loss.

Ex. 2. If $5\frac{1}{2}$ per cent. be gained by selling butter at £5. 5s. 6d. per cwt., what will be the gain per cent., by selling it at 1s. 3d. per lb.?

$$£5. 5s. 6d. = £5\frac{11}{40} = £\frac{211}{40}.$$

$$\begin{array}{ccc} £. & £. & £. \\ 105\frac{1}{2} : \frac{211}{40} :: 100 : \text{cost price;} \end{array}$$

$$\begin{aligned} \therefore \text{cost price} &= £ \frac{\cancel{211} \times \cancel{100} \times 2}{\cancel{40} \times \cancel{105} \times 2} \\ &= £5. \end{aligned}$$

$$\left(£105\frac{1}{2} \text{ and } £\frac{211}{40} \text{ are both selling prices.} \right)$$

If 1 lb. is sold for 1s. 3d.;

$$\therefore 1 \text{ cwt. is sold for } \frac{5}{4} \times \frac{\cancel{28}}{\cancel{1}} \text{ s.} = 14\text{os.} = £7;$$

$$\therefore \text{gain on 1 cwt.} = £7 - £5 = £2.$$

$$\begin{array}{ccc} £. & £. & £. \\ 5 : 100 :: 2 : \text{gain per cent.} \end{array}$$

$$\therefore \text{Ans.} = \frac{2 \times \cancel{100}}{\cancel{5}} \text{ per cent.}$$

$$= 40 \text{ per cent.}$$

N.B. Here £5 and £100 are both *cost prices*.

Ex. 3. If a person purchase pins when they are 18 in a row, and sell them 11 in a row at the same price, how much is his gain per cent. ?

Suppose 18 pins to cost 18 pence ;

\therefore 11 pins are sold for 18 pence.

But since 18 pins cost 18 pence ;

\therefore 11 pins cost 11 pence ;

\therefore 11 pins cost 11 pence, and are sold for 18 pence ;

\therefore 7 pence is the gain on the cost price 11 pence.

$$\begin{array}{ccc} d. & d. & d. \\ 11 : 100 :: 7 : \text{Ans.} \end{array}$$

N.B. 11*d.* and 100*d.* are both cost prices ; 7*d.* and the answer are both gains.

Art. 42. If two fruit-women were willing to exchange 12 apples for 49 pears, 121 nuts for 25 pears, 6 oranges for 5 plums, 154 nuts for 45 oranges ; how many apples would they be willing to exchange for 77 plums ?

$$\begin{array}{ccc} \text{pl.} & \text{pl.} & \text{or.} \\ 5 : 77 :: 6 : \text{no. of oranges ;} \end{array}$$

$$\therefore 77 \text{ plums are equivalent to } \frac{6 \times 77}{5} \text{ oranges.}$$

$$\begin{array}{ccc} \text{or.} & \text{or.} & \text{nuts.} \\ 45 : \frac{6 \times 77}{5} :: 154 : \text{no. of nuts.} \end{array}$$

$$\therefore 77 \text{ plums } \left(= \frac{6 \times 77}{5} \text{ oranges} \right) \text{ are equivalent to}$$

$$\frac{6 \times 77 \times 154}{5 \times 45} \text{ nuts.}$$

divisible by 9, that is, if the sum of the digits is divisible by 9. This agrees with the rule.

A similar proof would apply to the number 3.

A number is divisible by 11, if the difference of the sums of the digits in the odd and even places is divisible by 11.

Take the number 865432.

$$\begin{aligned}
 865432 &= 800000 + 60000 + 5000 + 400 + 30 + 2 \\
 &= 2 + 3(11 - 1) + 4(99 + 1) + 5(1001 - 1) + 6(9999 + 1) \\
 &\quad + 8(100001 - 1) \\
 &= (2 - 3 + 4 - 5 + 6 - 8) + (3 \times 11 + 4 \times 99 + 5 \times 1001 \\
 &\quad + 6 \times 9999 + 8 \times 100001).
 \end{aligned}$$

Now the second bracket is divisible by 11 (because 11, 99, 1001, 9999, 100001, are each divisible by 11); therefore the whole number is divisible by 11, if the first bracket is divisible by 11, that is, if the difference of the sums of the digits in the odd and even places is divisible by 11. This agrees with the rule.

MACMILLAN AND CO'S
PUBLICATIONS.

Works by BARNARD SMITH, M.A. *Rector of Glaston,
Rutland, &c.*

ARITHMETIC AND ALGEBRA in their Principles
and Application. Tenth Edition. Crown 8vo., 10s. 6d.

ARITHMETIC FOR THE USE OF SCHOOLS.
Ninth Edition. Crown 8vo., 4s. 6d. A KEY. Seventh Edition.
Crown 8vo., 8s. 6d.

EXERCISES IN ARITHMETIC. With Answers. Cr.
8vo., limp cloth, 2s. 6d. Or sold separately as follows :—Part I.
1s. Part II. 1s. Answers, 6d.

SCHOOL CLASS BOOK OF ARITHMETIC. 18mo.
3s. Or sold separately, Parts I. and II. 10d. Part III. 1s. KEY.
Complete in One Volume, 18mo., 6s. 6d.; or Parts I. II. and III.
2s. 6d. each.

SHILLING BOOK OF ARITHMETIC FOR NA-
TIONAL AND ELEMENTARY SCHOOLS. 18mo., cloth.
Or separately, Part I. 2d.; II. 3d.; III. 7d. With Answers, 1s. 6d.

ANSWERS TO THE SHILLING BOOK OF ARITH-
METIC. 18mo., 6d. KEY, 18mo., 4s. 6d.

EXAMINATION PAPERS IN ARITHMETIC. In
Four Parts. 18mo., 1s. 6d. With Answers, 1s. 9d. KEY, 18mo.,
4s. 6d.

ELEMENTARY MENSURATION. For the use of
Schools. With numerous Examples. By SEPTIMUS TEBAY, B.A.,
Head Master of Queen Elizabeth's Grammar School, Rivington.
Extra fcap. 8vo., 3s. 6d.

A GEOMETRICAL NOTE-BOOK. Containing Easy
Problems in Geometrical Drawing, preparatory to the study of
Geometry. For the use of Schools. By F. E. KITCHENER,
M.A., Mathematical Master at Rugby. 4to., 2s.

MODERN METHODS IN ELEMENTARY GEO-
METRY. By E. M. REYNOLDS, M.A., Mathematical Master
in Clifton College. Crown 8vo., 3s. 6d.

Works by ISAAC TODHUNTER, M.A., F.R.S.

EUCLID FOR COLLEGES AND SCHOOLS. New Edition. 18mo., 3s. 6d.

ALGEBRA FOR BEGINNERS. With numerous Examples. New Edition. 18mo., 2s. 6d. KEY. Crown 8vo., 6s. 6d.

MECHANICS FOR BEGINNERS. With numerous Examples. 18mo., 4s. 6d.

TRIGONOMETRY FOR BEGINNERS. With numerous Examples. 18mo., 2s. 6d.

ALGEBRA FOR THE USE OF COLLEGES AND SCHOOLS. Fourth Edition. Crown 8vo., 7s. 6d.

PLANE TRIGONOMETRY FOR COLLEGES AND SCHOOLS. Third Edition. Crown 8vo., 5s.

A TREATISE ON SPHERICAL TRIGONOMETRY FOR THE USE OF COLLEGES AND SCHOOLS. Second Edition. Crown 8vo., 4s. 6d.

ARITHMETICAL EXAMPLES. Progressively Arranged, with Exercises and Examination Papers. By the Rev. T. DALTON, M.A., Assistant-Master at Eton College. 18mo., 2s. 6d.

ALGEBRAICAL EXERCISES. Progressively Arranged. By the Rev. C. A. JONES, M.A., and C. H. CHEYNE, M.A. New Edition. 18mo., 2s. 6d.

ELEMENTARY GEOMETRY. Part I. Angles, Parallels, Triangles, &c. Compiled by J. M. WILSON, M.A., Mathematical Master of Rugby School. Extra fcap. 8vo., 2s. 6d.

ELEMENTARY GEOMETRY. Part II. The Circle and Proportion. By J. M. WILSON, M.A. Extra fcap. 8vo., 2s. 6d.

CLASSICS.

SALLUST. With English Notes. By C. MERIVALE, B.D., Author of "History of Rome," &c. Second Edition. Fcp. 8vo., 4s. 6d.

The **CATILINA** and **JUGURTHA** may be had separately, price 2s. 6d. each, bound in cloth.

CICERO'S SECOND PHILIPPIC. With English Notes. By JOHN E. B. MAYOR. Third Edition. Fcap. 8vo., 5s.

THE SICILIAN EXPEDITION, being Books VI. and VII. of Thucydides, with Notes. By the Rev. PERCIVAL FROST, M.A., late Fellow of St. John's College, Cambridge. A new Edition, revised and enlarged, with a Map. Extra fcap. 8vo., 5s.

DEMOSTHENES ON THE CROWN. With English Notes. By B. DRAKE, M.A. Third Edition. To which is prefixed, **ÆSCHINES** against **CTESIPHON**. With English Notes. Fcp. 8vo., 5s.

MYTHOLOGY FOR LATIN VERSIFICATION. A Brief Sketch of the Fables of the Ancients, prepared to be rendered into Latin Verse for Schools. By F. HODGSON, B.D. New Edition, Revised by F. C. HODGSON, M.A. 18mo., 3s.

TACITUS; THE AGRICOLA AND GERMANY. Translated into English by A. J. CHURCH and W. J. BRODRIBB. With Map and Notes. Fcap. 8vo., 2s. 6d.

FIRST GREEK READER. Edited after KARL HALM. With Additions and Corrections. By J. E. B. MAYOR, M.A. Fcp. 8vo., 6s.

Works by J. WRIGHT, M.A.

HELLENICA; OR, A HISTORY OF GREECE IN Greek, being a First Greek Reading Book, with Explanatory Notes. *Third Edition.* WITH A VOCABULARY. 12mo., 3s. 6d.

THE SEVEN KINGS OF ROME. An Easy Narrative, abridged from the First Book of Livy by the omission of difficult passages, being a First Latin Reading Book, with Grammatical Notes. Fcap. 8vo., 3s.

A VOCABULARY AND EXERCISES ON THE 'SEVEN Kings of Rome.' Fcap. 8vo., 2s. 6d.

*• The Vocabulary and Exercises may also be had bound up with 'The Seven Kings of Rome.' Price 5s.

A HELP TO LATIN GRAMMAR; Or, the Form and Use of Words in Latin, with Progressive Exercises. Crown 8vo., 4s. 6d.

ELEMENTARY SCIENCE.

LESSONS IN ELEMENTARY PHYSIOLOGY. With numerous Illustrations. By T. H. HUXLEY, F.R.S., Professor of Natural History in the Royal School of Mines. Fifth Thousand. 18mo., 4s. 6d.

QUESTIONS ON HUXLEY'S PHYSIOLOGY. For Schools. By J. ALCOCK. 18mo. 1s. 6d.

POPULAR ASTRONOMY. With Illustrations. By G. B. AIRY, Astronomer Royal. Sixth and Cheaper Edition. 18mo. 4s. 6d.

LESSONS IN ELEMENTARY CHEMISTRY. By HENRY ROSCOE, F.R.S., Professor of Chemistry in Owen's College, Manchester. With numerous Illustrations and Chromo.-Litho. of the Solar Spectra. Twelfth Thousand. 18mo., 4s. 6d.

LESSONS IN ELEMENTARY BOTANY. With nearly 200 Illustrations. By DANIEL OLIVER, F.R.S., F.L.S. Fourth Thousand. 18mo., 4s. 6d.

ELEMENTARY LESSONS IN ASTRONOMY. By J. NORMAN LOCKYER, F.R.A.S. With Coloured Diagram and numerous Woodcuts. 18mo., 5s. 6d.

Works by the REV. G. F. MACLEAR, B.D., *Head Master of King's College School.*

A **CLASS-BOOK OF OLD TESTAMENT HISTORY.** Fourth Edition, with Four Maps. 18mo., 4s. 6d.

A **CLASS-BOOK OF NEW TESTAMENT HISTORY:** including the Connection of the Old and New Testament. With Four Maps. Second Edition. 18mo., 5s. 6d.

A **CLASS-BOOK OF THE CATECHISM OF THE CHURCH OF ENGLAND.** 18mo., 2s. 6d.

A **SHILLING BOOK OF OLD TESTAMENT HISTORY,** for National and Elementary Schools. With Map. 18mo., cloth limp.

A **SHILLING BOOK OF NEW TESTAMENT HISTORY,** for National and Elementary Schools. With Map. 18mo., cloth limp.

A **FIRST CLASS BOOK OF THE CATECHISM OF THE CHURCH OF ENGLAND,** with Scripture Proofs, for Junior Classes and Schools. 18mo., 6d.

NOVEMBER, 1868.

LIST OF EDUCATIONAL BOOKS

PUBLISHED BY

MACMILLAN AND CO.,

16, *BEDFORD STREET, COVENT GARDEN,*

London, w.c.

.....

CONTENTS.

	Page
CLASSICAL	3
MATHEMATICAL	7
SCIENCE	17
MISCELLANEOUS	19
DIVINITY	21
BOOKS ON EDUCATION	24

MESSRS. MACMILLAN & CO. beg to call attention to the accompanying Catalogue of their EDUCATIONAL WORKS, the writers of which are mostly scholars of eminence in the Universities, as well as of large experience in teaching.

Many of the works have already attained a wide circulation in England and in the Colonies, and are acknowledged to be among the very best Educational Books on their respective subjects.

The books can generally be procured by ordering them through local booksellers in town or country, but if at any time difficulty should arise, Messrs. MACMILLAN will feel much obliged by direct communication with themselves on the subject.

Notices of errors or defects in any of these works will be gratefully received and acknowledged.

LIST OF EDUCATIONAL BOOKS.

CLASSICAL.

ÆSCHYLI EUMENIDES. The Greek Text, with English Notes, and English Verse Translation and an Introduction. By **BERNARD DRAKE**, M.A., late Fellow of King's College, Cambridge. 8vo. 7s. 6d.

The Greek Text adopted in this Edition is based upon that of Wellauer, which may be said in general terms to represent that of the best manuscripts. But in correcting the Text, and in the Notes, advantage has been taken of the suggestions of Hermann, Paley, Linwood, and other commentators.

ARISTOTLE ON FALLACIES; OR, THE SOPHISTICI ELENCHI. With a Translation and Notes by **EDWARD POSTE**, M.A., Fellow of Oriel College, Oxford. 8vo. 8s. 6d.

Besides the doctrine of Fallacies, Aristotle offers either in this treatise, or in other passages quoted in the commentary, various glances over the world of science and opinion, various suggestions on problems which are still agitated, and a vivid picture of the ancient system of dialectics, which it is hoped may be found both interesting and instructive.

"It is not only scholarlike and careful; it is also perspicuous."—*Guardian*.

ARISTOTLE.—AN INTRODUCTION TO ARISTOTLE'S RHETORIC. With Analysis, Notes, and Appendices. By **E. M. COPE**, Senior Fellow and Tutor of Trinity College, Cambridge. 8vo. 14s.

This work is introductory to an edition of the Greek Text of Aristotle's Rhetoric, which is in course of preparation.

"Mr. Cope has given a very useful appendage to the promised Greek Text; but also a work of so much independent use that he is quite justified in his separate publication. All who have the Greek Text will find themselves supplied with a comment; and those who have not will find an analysis of the work."—*Athenæum*.

CATULLI VERONENSIS LIBER, edited by R. ELLIS, Fellow of Trinity College, Oxford. 18mo. 3s. 6d.

"It is little to say that no edition of Catullus at once so scholarlike has ever appeared in England."—*Athenaeum*.

"Rarely have we read a classic author with so reliable, acute, and safe a guide."—*Saturday Review*.

CICERO.—THE SECOND PHILIPPIC ORATION. With an Introduction and Notes, translated from the German of KARL HAEM. Edited, with Corrections and Additions, by JOHN E. B. MAYOR, M.A., Fellow and Classical Lecturer of St. John's College, Cambridge. Third Edition, revised. Fcap. 8vo. 5s.

"A very valuable edition, from which the student may gather much both in the way of information directly communicated, and directions to other sources of knowledge."—*Athenaeum*.

DEMOSTHENES ON THE CROWN. The Greek Text with English Notes. By B. DRAKE, M.A., late Fellow of King's College, Cambridge. Third Edition, to which is prefixed ÆSCHINES AGAINST CTESIPHON, with English Notes. Fcap. 8vo. 5s.

The terseness and felicity of Mr. Drake's translations constitute perhaps the chief value of his edition, and the historical and archæological details necessary to understanding the *De Coroniâ* have in some measure been anticipated in the notes on the Oration of Æschines. In both, the text adopted in the Zurich edition of 1851, and taken from the Parisian MS., has been adhered to without any variation. Where the readings of Bekker, Dissen, and others appear preferable, they are subjoined in the notes.

HODGSON.—MYTHOLOGY FOR LATIN VERSIFICATION. A Brief Sketch of the Fables of the Ancients, prepared to be rendered into Latin Verse for Schools. By F. HODGSON, B.D., late Provost of Eton. New Edition, revised by F. C. HODGSON, M.A. 18mo. 3s.

Intending the little book to be entirely elementary, the Author has made it as easy as he could, without too largely superseding the use of the Dictionary and Gradus. By the facilities here afforded, it will be possible, in many cases, for a boy to get rapidly through these preparatory exercises; and thus, having mastered the first difficulties, he may advance with better hopes of improvement to subjects of higher character, and verses of more difficult composition.

JUVENAL, FOR SCHOOLS. With English Notes. By J. E. B. MAYOR, M.A. New and Cheaper Edition. Crown 8vo.

[*In the Press.*]

"A School edition of Juvenal, which, for really ripe scholarship, extensive acquaintance with Latin literature, and familiar knowledge of Continental criticism, ancient and modern, is unsurpassed, we do not say among English School-books, but among English editions generally."—*Edinburgh Review.*

LYTTELTON.—THE COMUS of MILTON rendered into Greek Verse. By LORD LYTTELTON. Extra fcap. 8vo. Second Edition. 5s.

— THE SAMSON AGONISTES of MILTON rendered into Greek Verse. By LORD LYTTELTON. Extra fcap. 8vo. 6s. 6d.

MARSHALL.—A TABLE OF IRREGULAR GREEK VERBS, Classified according to the Arrangement of Curtius's Greek Grammar. By J. M. MARSHALL, M.A., Fellow and late Lecturer of Brasenose College, Oxford; one of the Masters in Clifton College. 8vo. cloth. 1s.

MAYOR.—FIRST GREEK READER. Edited after KARL HALM, with Corrections and large Additions by JOHN E. B. MAYOR, M.A., Fellow and Classical Lecturer of St. John's College, Cambridge. Fcap. 8vo. 6s.

MERIVALE.—KEATS' HYPERION rendered into Latin Verse. By C. MERIVALE, B.D. Second Edition. Extra fcap. 8vo. 3s. 6d.

PLATO.—THE REPUBLIC OF PLATO. Translated into English, with an Analysis and Notes, by J. LL. DAVIES, M.A., and D. J. VAUGHAN, M.A. Third Edition, with Vignette Portraits of Plato and Socrates, engraved by JEENS from an Antique Gem. 18mo. 4s. 6d.

ROBY.—A LATIN GRAMMAR for the Higher Classes in Grammar Schools. By H. J. ROBY, M.A.; based on the "Elementary Latin Grammar." [*In the Press.*]

SALLUST.—**C. CAII SALLUSTII CRISPI** *Catilina et Jugurtha*. For use in Schools (with copious Notes). By C. MERIVALE, B.D. (In the present Edition the Notes have been carefully revised, and a few remarks and explanations added.) Second Edition. Fcap. 8vo. 4s. 6d.

The *Jugurtha* and the *Catilina* may be had separately, price 2s. 6d. each.

TACITUS.—**THE HISTORY OF TACITUS** translated into ENGLISH. By A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A. With Notes and a Map. 8vo. 10s. 6d.

The translators have endeavoured to adhere as closely to the original as was thought consistent with a proper observance of English idiom. At the same time it has been their aim to reproduce the precise expressions of the author. The campaign of *Civilis* is elucidated in a note of some length which is illustrated by a map, containing only the names of places and of tribes occurring in the work.

— **THE AGRICOLA and GERMANY.** By the same translators. With Maps and Notes. Extra fcap. 8vo. 2s. 6d.

THRING.—Works by **Edward Thring, M.A.**, Head Master of Uppingham School:—

— **A CONSTRUING BOOK.** Fcap. 8vo. 2s. 6d.

This Construing Book is drawn up on the same sort of graduated scale as the Author's *English Grammar*. Passages out of the best Latin Poets are gradually built up into their perfect shape. The few words altered, or inserted as the passages go on, are printed in Italics. It is hoped by this plan that the learner, whilst acquiring the rudiments of language, may store his mind with good poetry and a good vocabulary.

— **A LATIN GRADUAL.** A First Latin Construing Book for Beginners. Fcap. 8vo. 2s. 6d.

The main plan of this little work has been well tested.

The intention is to supply by easy steps a knowledge of Grammar, combined with a good vocabulary; in a word, a book which will not require to be forgotten again as the learner advances.

A short practical manual of common Mood constructions, with their English equivalents, form the second part.

— **A MANUAL of MOOD CONSTRUCTIONS.** Extra fcap. 8vo. 1s. 6d.

THUCYDIDES.—**THE SICILIAN EXPEDITION.** Being Books VI. and VII. of Thucydides, with Notes. A New Edition, revised and enlarged, with a Map. By the Rev. PERCIVAL FROST, M.A., late Fellow of St. John's College, Cambridge. Fcap. 8vo. 5s.

This edition is mainly a grammatical one. Attention is called to the force of compound verbs, and the exact meaning of the various tenses employed.

WRIGHT.—Works by **J. Wright, M.A.**, late Head Master of Sutton Coldfield School:—

- **HELLENICA**; Or, a **HISTORY of GREECE in GREEK**, as related by Diodorus and Thucydides, being a First Greek Reading Book, with Explanatory Notes Critical and Historical. Second Edition, with a Vocabulary. 12mo. 3s. 6d.

In the last twenty chapters of this volume, Thucydides sketches the rise and progress of the Athenian Empire in so clear a style and in such simple language, that the author doubts whether any easier or more instructive passages can be selected for the use of the pupil who is commencing Greek.

- **A HELP TO LATIN GRAMMAR**; Or, the Form and Use of Words in Latin, with Progressive Exercises. Crown 8vo. 4s. 6d.

"Never was there a better aid offered alike to teacher and scholar in that arduous pass. The style is at once familiar and strikingly simple and lucid; and the explanations precisely hit the difficulties, and thoroughly explain them."—*English Journal of Education*.

- **THE SEVEN KINGS OF ROME**. An Easy Narrative, abridged from the First Book of Livy by the omission of difficult passages, being a First Latin Reading Book, with Grammatical Notes. Fcap. 8vo. 3s.

This work is intended to supply the pupil with an easy Construing-book, which may at the same time be made the vehicle for instructing him in the rules of grammar and principles of composition. Here Livy tells his own pleasant stories in his own pleasant words. Let Livy be the master to teach a boy Latin, not some English collector of sentences, and he will not be found a dull one.

- **A VOCABULARY AND EXERCISES** on the "**SEVEN KINGS OF ROME**." Fcap. 8vo. 2s. 6d.

The Vocabulary and Exercises may also be had bound up with "**The Seven Kings of Rome**," price 5s.

MATHEMATICAL.

AIRY.—Works by **G. B. Airy**, Astronomer Royal:—

- **ELEMENTARY TREATISE ON PARTIAL DIFFERENTIAL EQUATIONS**. Designed for the use of Students in the University. With Diagrams. Crown 8vo. cloth, 5s. 6d.

It is hoped that the methods of solution here explained, and the instances exhibited, will be found sufficient for application to nearly all the important problems of Physical Science, which require for their complete investigation the aid of partial differential equations.

AIRY.—Works by **G. B. Airy**—*Continued*.

- ON THE ALGEBRAICAL AND NUMERICAL THEORY of ERRORS of OBSERVATIONS, and the COMBINATION of OBSERVATIONS. Crown 8vo. cloth, 6s. 6d.
- UNDULATORY THEORY OF OPTICS. Designed for the use of Students in the University. New Edition. Crown 8vo. cloth, 6s. 6d.
- ON SOUND and ATMOSPHERIC VIBRATIONS. With the Mathematical Elements of Music. Designed for the use of Students of the University. Crown 8vo. 9s.

BAYMA.—THE ELEMENTS of MOLECULAR MECHANICS.

By JOSEPH BAYMA, S.J., Professor of Philosophy, Stonyhurst College. Demy 8vo. cloth, 10s. 6d.

BOOLE.—Works by **G. Boole, D.C.L., F.R.S.**, Professor of Mathematics in the Queen's University, Ireland :—

- A TREATISE ON DIFFERENTIAL EQUATIONS. New and Revised Edition. Edited by I. TODHUNTER. Crown 8vo. cloth, 14s.

The author has endeavoured in this Treatise to convey as complete an account of the present state of knowledge on the subject of Differential Equations, as was consistent with the idea of a work intended primarily for elementary instruction. The earlier sections of each chapter contain that kind of matter which has usually been thought suitable to the beginner, while the later ones are devoted either to an account of recent discovery, or the discussion of such deeper questions of principle as are likely to present themselves to the reflective student in connexion with the methods and processes of his previous course.

- A TREATISE ON DIFFERENTIAL EQUATIONS. Supplementary Volume. Edited by I. TODHUNTER. Crown 8vo. cloth, 8s. 6d.
- THE CALCULUS OF FINITE DIFFERENCES. Crown 8vo. cloth, 10s. 6d.

This work is in some measure designed as a sequel to the *Treatise on Differential Equations*, and is composed on the same plan.

BEASLEY.—AN ELEMENTARY TREATISE ON PLANE TRIGONOMETRY. With Examples. By R. D. BEASLEY, M.A., Head Master of Grantham Grammar School. Second Edition, revised and enlarged. Crown 8vo. cloth, 3s. 6d.

This Treatise is specially intended for use in Schools. The choice of matter has been chiefly guided by the requirements of the three days' Examination at Cambridge, with the exception of proportional parts in Logarithms, which have been omitted. About *Four hundred* Examples have been added, mainly collected from the Examination Papers of the last ten years, and great pains have been taken to exclude from the body of the work any which might dishearten a beginner by their difficulty.

CAMBRIDGE SENATE-HOUSE PROBLEMS and RIDERS, WITH SOLUTIONS:—

1848—1851.—PROBLEMS. By FERRERS and JACKSON. 8vo. cloth. 15s. 6d.

1848—1851.—RIDERS. By JAMESON. 8vo. cloth. 7s. 6d.

1854.—PROBLEMS and RIDERS. By WALTON and MACKENZIE, 8vo. cloth. 10s. 6d.

1857.—PROBLEMS and RIDERS. By CAMPION and WALTON. 8vo. cloth. 8s. 6d.

1860.—PROBLEMS and RIDERS. By WATSON and ROUTH. Crown 8vo. cloth. 7s. 6d.

1864.—PROBLEMS and RIDERS. By WALTON and WILKINSON. 8vo. cloth. 10s. 6d.

CAMBRIDGE COURSE OF ELEMENTARY NATURAL PHILOSOPHY, for the Degree of B.A. Originally compiled by J. C. SNOWBALL, M.A., late Fellow of St. John's College. Fifth Edition, revised and enlarged, and adapted for the Middle-Class Examinations by THOMAS LUND, B.D., Late Fellow and Lecturer of St. John's College; Editor of Wood's Algebra, &c. Crown 8vo. cloth. 5s.

This work will be found suited to the wants, not only of University Students, but also of many others who require a short course of Mechanics and Hydrostatics, and especially of the Candidates at our Middle-Class Examinations.

CAMBRIDGE AND DUBLIN MATHEMATICAL JOURNAL.

THE COMPLETE WORK, in Nine Vols. 8vo. cloth. £7 4s.
(Only a few copies remain on hand.)

CHEYNE.—AN ELEMENTARY TREATISE on the PLANETARY THEORY. With a Collection of Problems. By C. H. H. CHEYNE, B.A. Crown 8vo. cloth. 6s. 6d.

— THE EARTH'S MOTION of ROTATION. By C. H. H. CHEYNE, M.A. Crown 8vo. 3s. 6d.

CHILDE.—THE SINGULAR PROPERTIES of the ELLIPSOID and ASSOCIATED SURFACES of the n th DEGREE. By the Rev. G. F. CHILDE, M.A., Author of "Ray Surfaces," "Related Caustics," &c. 8vo. 10s. 6d.

CHRISTIE.—A COLLECTION OF ELEMENTARY TEST-QUESTIONS in PURE and MIXED MATHEMATICS; with Answers and Appendices on Synthetic Division, and on the Solution of Numerical Equations by Horner's Method. By JAMES R. CHRISTIE, F.R.S., late First Mathematical Master at the Royal Military Academy, Woolwich. Crown 8vo. cloth, 8s. 6d.

DALTON.—ARITHMETICAL EXAMPLES. Progressively arranged, with Exercises and Examination Papers. By the Rev. T. DALTON, M.A., Assistant Master of Eton College. 18mo. cloth. 2s. 6d.

DAY.—PROPERTIES OF CONIC SECTIONS PROVED GEOMETRICALLY. Part I., THE ELLIPSE, with Problems. By the Rev. H. G. DAY, M.A., Head Master of Sedbergh Grammar School. Crown 8vo. 3s. 6d.

DODGSON.—AN ELEMENTARY TREATISE ON DETERMINANTS, with their Application to Simultaneous Linear Equations and Algebraical Geometry. By C. L. DODGSON, M.A., Mathematical Lecturer of Christ Church, Oxford. Small 4to. cloth, 10s. 6d.

DREW.—GEOMETRICAL TREATISE on CONIC SECTIONS. By W. H. DREW, M.A., St. John's College, Cambridge. Third Edition. Crown 8vo. cloth, 4s. 6d.

In this work the subject of Conic Sections has been placed before the student in such a form that, it is hoped, after mastering the elements of Euclid, he may find it an easy and interesting continuation of his geometrical studies. With a view also of rendering the work a complete Manual of what is required at the Universities, there have been either embodied into the text, or inserted among the examples, every book-work question, problem, and rider, which has been proposed in the Cambridge examinations up to the present time.

— SOLUTIONS TO THE PROBLEMS IN DREW'S CONIC SECTIONS. Crown 8vo. cloth, 4s. 6d.

FERRERS.—AN ELEMENTARY TREATISE on TRILINEAR CO-ORDINATES, the Method of Reciprocal Polars, and the Theory of Projections. By the Rev. N. M. FERRERS, M.A., Fellow and Tutor of Gonville and Caius College, Cambridge. Second Edition. Crown 8vo. 6s. 6d.

The object of the author in writing on this subject has mainly been to place it on a basis altogether independent of the ordinary Cartesian system, instead of regarding it as only a special form of Abridged Notation. A short chapter on Determinants has been introduced.

FROST.—THE FIRST THREE SECTIONS of NEWTON'S PRINCIPIA. With Notes and Illustrations. Also a Collection of Problems, principally intended as Examples of Newton's Methods. By PERCIVAL FROST, M.A., late Fellow of St. John's College, Mathematical Lecturer of King's College, Cambridge. Second Edition. 8vo. cloth, 10s. 6d.

The author's principal intention is to explain difficulties which may be encountered by the student on first reading the Principia, and to illustrate the advantages of a careful study of the methods employed by Newton, by showing the extent to which they may be applied in the solution of problems; he has also endeavoured to give assistance to the student who is engaged in the study of the higher branches of Mathematics, by representing in a geometrical form several of the processes employed in the Differential and Integral Calculus, and in the analytical investigations of Dynamics.

FROST and WOLSTENHOLME.—A TREATISE ON SOLID GEOMETRY. By PERCIVAL FROST, M.A., and the Rev. J. WOLSTENHOLME, M.A., Fellow and Assistant Tutor of Christ's College. 8vo. cloth, 18s.

The authors have endeavoured to present before students as comprehensive a view of the subject as possible. Intending as they have done to make the subject accessible, at least in the earlier portion, to all classes of students, they have endeavoured to explain fully all the processes which are most useful in dealing with ordinary theorems and problems, thus directing the student to the selection of methods which are best adapted to the exigencies of each problem. In the more difficult portions of the subject, they have considered themselves to be addressing a higher class of students; there they have tried to lay a good foundation on which to build, if any reader should wish to pursue the science beyond the limits to which the work extends.

GODFRAY.—A TREATISE on ASTRONOMY, for the use of Colleges and Schools. By HUGH GODFRAY, M.A., Mathematical Lecturer at Pembroke College, Cambridge. 8vo. cloth. 12s. 6d.

"We can recommend for its purpose a very good *Treatise on Astronomy* by Mr. Godfray. It is a working book, taking astronomy in its proper place in mathematical science. But it begins with the elementary definitions, and connects the mathematical formulæ very clearly with the visible aspect of the heavens and the instruments which are used for observing it."—*Guardian*.

— AN ELEMENTARY TREATISE on the LUNAR THEORY. With a brief Sketch of the Problem up to the time of Newton. By HUGH GODFRAY, M.A. Second Edition, revised. Crown 8vo. cloth. 5s. 6d.

HEMMING.—AN ELEMENTARY TREATISE on the DIFFERENTIAL AND INTEGRAL CALCULUS, for the use of Colleges and Schools. By G. W. HEMMING, M.A., Fellow of St. John's College, Cambridge. Second Edition, with Corrections and Additions. 8vo. cloth. 9s.

JONES and CHEYNE.—ALGEBRAICAL EXERCISES. Progressively arranged. By the Rev. C. A. JONES, M.A., and C. H. CHEYNE, M.A., Mathematical Masters of Westminster School. New Edition. 18mo. cloth, 2s. 6d.

This little book is intended to meet a difficulty which is probably felt more or less by all engaged in teaching Algebra to beginners. It is that while new ideas are being acquired, old ones are forgotten. In the belief that constant practice is the only remedy for this, the present series of miscellaneous exercises has been prepared. Their peculiarity consists in this, that though miscellaneous they are yet progressive, and may be used by the pupil almost from the commencement of his studies. They are not intended to supersede the systematically arranged examples to be found in ordinary treatises on Algebra, but rather to supplement them.

The book being intended chiefly for Schools and Junior Students, the higher parts of Algebra have not been included.

KITCHENER.—A GEOMETRICAL NOTE-BOOK, containing Easy Problems in Geometrical Drawing preparatory to the Study of Geometry. For the use of Schools. By F. E. KITCHENER, M.A., Mathematical Master at Rugby. 4to. 2s.

MORGAN.—A COLLECTION of PROBLEMS and EXAMPLES in Mathematics. With Answers. By H. A. MORGAN, M.A., Sadlerian and Mathematical Lecturer of Jesus College, Cambridge. Crown 8vo. cloth. 6s. 6d.

This book contains a number of problems, chiefly elementary, in the Mathematical subjects usually read at Cambridge. They have been selected from the papers set during late years at Jesus college. Very few of them are to be met with in other collections, and by far the larger number are due to some of the most distinguished Mathematicians in the University.

PARKINSON.—Works by **S. Parkinson, B.D.**, Fellow and Prælector of St. John's College, Cambridge :—

- **AN ELEMENTARY TREATISE ON MECHANICS.** For the use of the Junior Classes at the University and the Higher Classes in Schools. With a Collection of Examples. Third Edition, revised. Crown 8vo. cloth, 9s. 6d.

The author has endeavoured to render the present volume suitable as a Manual for the junior classes in Universities and the higher classes in Schools. In the Third Edition several additional propositions have been incorporated in the work for the purpose of rendering it more complete, and the Collection of Examples and Problems has been largely increased.

- **A TREATISE ON OPTICS.** Second Edition, revised. Crown 8vo. cloth, 10s. 6d.

A collection of Examples and Problems has been appended to this work which are sufficiently numerous and varied in character to afford useful exercise for the student: for the greater part of them recourse has been had to the Examination Papers set in the University and the several Colleges during the last twenty years.

PHEAR.—ELEMENTARY HYDROSTATICS. With numerous Examples. By J. B. PHEAR, M.A., Fellow and late Assistant Tutor of Clare College, Cambridge. Fourth Edition. Crown 8vo. cloth, 5s. 6d.

"An excellent Introductory Book. The definitions are very clear; the descriptions and explanations are sufficiently full and intelligible; the investigations are simple and scientific. The examples greatly enhance its value."—*English Journal of Education*.

PRATT.—A TREATISE on ATTRACTIONS, LAPLACE'S FUNCTIONS, and the FIGURE of the EARTH. By JOHN H. PRATT, M.A., Archdeacon of Calcutta, Author of "The Mathematical Principles of Mechanical Philosophy." Third Edition. Crown 8vo. cloth, 6s. 6d.

PUCKLE.—AN ELEMENTARY TREATISE on CONIC SECTIONS and ALGEBRAIC GEOMETRY. With numerous Examples and hints for their Solution; especially designed for the use of Beginners. By G. H. PUCKLE, M.A., St. John's College, Cambridge, Head Master of Windermere College. Third Edition, enlarged and improved. Crown 8vo. cloth, 7s. 6d.

The work has been completely re-written, and a considerable amount of new matter has been added, to suit the requirements of the present time.

RAWLINSON.—ELEMENTARY STATICS. By G. RAWLINSON, M.A. Edited by EDWARD STURGES, M.A., of Emmanuel College, Cambridge, and late Professor of the Applied Sciences, Elphinstone College, Bombay. Crown 8vo. cloth. 4s. 6d.

Published under the authority of H. M. Secretary of State for use in the Government Schools and Colleges in India.

"This Manual may take its place among the most exhaustive, yet clear and simple, we have met with, upon the composition and resolution of forces, equilibrium, and the mechanical powers."—*Oriental Budget*.

REYNOLDS.—MODERN METHODS IN ELEMENTARY GEOMETRY. By E. M. REYNOLDS, M.A., Mathematical Master in Clifton College. Crown 8vo. 3s. 6d.

ROUTH.—AN ELEMENTARY TREATISE on the DYNAMICS of a SYSTEM of RIGID BODIES. With Examples. By EDWARD JOHN ROUTH, M.A., Fellow and Assistant Tutor of St. Peter's College, Cambridge; Examiner in the University of London. Crown 8vo. cloth, 10s. 6d.

SMITH.—A TREATISE on ELEMENTARY STATICS. By J. H. SMITH, M.A., Gonville and Caius College, Cambridge. 8vo. 5s. 6d.

SMITH.—Works by **Barnard Smith, M.A.**, Rector of Glaston, Rutlandshire, late Fellow and Senior Bursar of St. Peter's College, Cambridge :—

- **ARITHMETIC and ALGEBRA**, in their Principles and Application, with numerous Systematically arranged Examples, taken from the Cambridge Examination Papers, with especial reference to the Ordinary Examination for B.A. Degree. Tenth Edition. Crown 8vo. cloth, 10s. 6d.

This work is now extensively used in *Schools and Colleges* both *at home* and in the *Colonies*. It has also been found of great service for students preparing for the MIDDLE-CLASS AND CIVIL AND MILITARY SERVICE EXAMINATIONS, from the care that has been taken to elucidate the *principles* of all the Rules.

- **ARITHMETIC FOR SCHOOLS.** New Edition. Crown 8vo. cloth, 4s. 6d.
- **COMPANION to ARITHMETIC for SCHOOLS.** [*Preparing.*]
- **A KEY to the ARITHMETIC for SCHOOLS.** Seventh Edition. Crown 8vo., cloth, 8s. 6d.
- **EXERCISES in ARITHMETIC.** With Answers. Crown 8vo. limp cloth, 2s. 6d. Or sold separately, as follows :—Part I. 1s.; Part II. 1s. ANSWERS, 6d.

These Exercises have been published in order to give the pupil examples in every rule of Arithmetic. The greater number have been carefully compiled from the latest University and School Examination Papers.

- **SCHOOL CLASS-BOOK of ARITHMETIC.** 18mo. cloth, 3s. Or sold separately, Parts I. and II. 10d. each; Part III. 1s.
- **KEYS to SCHOOL CLASS-BOOK of ARITHMETIC.** Complete in one Volume, 18mo., cloth, 6s. 6d.; or Parts I., II., and III. 2s. 6d. each.
- **SHILLING BOOK of ARITHMETIC for NATIONAL and ELEMENTARY SCHOOLS.** 18mo. cloth. Or separately, Part I. 2d.; Part II. 3d.; Part III. 7d. ANSWERS, 6d.
- **THE SAME, with Answers complete.** 18mo. cloth, 1s. 6d.
- **KEY to SHILLING BOOK of ARITHMETIC.** 18mo. cloth, 4s. 6d.
- **EXAMINATION PAPERS in ARITHMETIC.** In Four Parts. 18mo. cloth, 1s. 6d. **THE SAME, with Answers,** 18mo. 1s. 9d.
- **KEY to EXAMINATION PAPERS in ARITHMETIC.** 18mo. cloth, 4s. 6d.

SNOWBALL.—PLANE and SPHERICAL TRIGONOMETRY. With the Construction and Use of Tables of Logarithms. By J. C. SNOWBALL. Tenth Edition. Crown 8vo. cloth, 7s. 6d.

TAIT and STEELE.—DYNAMICS of a PARTICLE. With Examples. By Professor TAIT and Mr. STEELE. New Edition. Crown 8vo. cloth, 10s. 6d.

In this Treatise will be found all the ordinary propositions connected with the Dynamics of Particles which can be conveniently deduced without the use of D'Alembert's Principles. Throughout the book will be found a number of illustrative Examples introduced in the text, and for the most part completely worked out; others, with occasional solutions or hints to assist the student, are appended to each Chapter.

TAYLOR.—GEOMETRICAL CONICS; including Anharmonic Ratio and Projection, with numerous Examples. By C. TAYLOR, B.A., Scholar of St. John's College, Cambridge. Crown 8vo. cloth, 7s. 6d.

TEBAY.—ELEMENTARY MENSURATION for SCHOOLS. With numerous Examples. By SEPTIMUS TEBAY, B.A., Head Master of Queen Elizabeth's Grammar School, Rivington. Extra fcap. 8vo. 3s. 6d.

TODHUNTER.—Works by I. Todhunter, M.A., F.R.S., Fellow and Principal Mathematical Lecturer of St. John's College, Cambridge:—

- THE ELEMENTS of EUCLID for the use of COLLEGES and SCHOOLS. New Edition. 18mo. cloth, 3s. 6d.
- ALGEBRA for BEGINNERS. With numerous Examples. New Edition. 18mo. cloth, 2s. 6d.
- KEY to ALGEBRA for BEGINNERS. Crown 8vo., cl., 6s. 6d.
- TRIGONOMETRY for BEGINNERS. With numerous Examples. New Edition. 18mo. cloth, 2s. 6d.

Intended to serve as an introduction to the larger treatise on *Plane Trigonometry*, published by the author. The same plan has been adopted as in the *Algebra for Beginners*: the subject is discussed in short chapters, and a collection of examples is attached to each chapter.

- MECHANICS for BEGINNERS. With numerous Examples. 18mo. cloth, 4s. 6d.

Intended as a companion to the two preceding books. The work forms an elementary treatise on *Demonstrative Mechanics*. It may be true that this part of mixed mathematics has been sometimes made too abstract and speculative; but it can hardly be doubted that a knowledge of the elements at least of the theory of the subject is extremely valuable even for those who are mainly concerned with practical results. The author has accordingly endeavoured to provide a suitable introduction to the study of applied as well as of theoretical Mechanics.

TODHUNTER.—Works by I. Todhunter, M.A.—*Continued.*

- A TREATISE on the DIFFERENTIAL CALCULUS. With Examples. Fourth Edition. Crown 8vo. cloth, 10s. 6d.
- A TREATISE on the INTEGRAL CALCULUS. Third Edition, revised and enlarged. With Examples. Crown 8vo. cloth, 10s. 6d.
- A TREATISE on ANALYTICAL STATICS. With Examples. Third Edition, revised and enlarged. Crown 8vo. cloth, 10s. 6d.
- PLANE CO-ORDINATE GEOMETRY, as applied to the Straight Line and the CONIC SECTIONS. With numerous Examples. Fourth Edition. Crown 8vo. cloth, 7s. 6d.
- ALGEBRA. For the use of Colleges and Schools. Fourth Edition. Crown 8vo. cloth, 7s. 6d.

This work contains all the propositions which are usually included in elementary treatises on Algebra, and a large number of *Examples for Exercise*. The author has sought to render the work easily intelligible to students without impairing the accuracy of the demonstrations, or contracting the limits of the subject. The Examples have been selected with a view to illustrate every part of the subject, and as the number of them is about *Sixteen hundred and fifty*, it is hoped they will supply ample exercise for the student. Each set of Examples has been carefully arranged, commencing with very simple exercises, and proceeding gradually to those which are less obvious.

- PLANE TRIGONOMETRY. For Schools and Colleges. Third Edition. Crown 8vo. cloth, 5s.

The design of this work has been to render the subject intelligible to beginners, and at the same time to afford the student the opportunity of obtaining all the information which he will require on this branch of Mathematics. Each chapter is followed by a set of Examples; those which are entitled *Miscellaneous Examples*, together with a few in some of the other sets, may be advantageously reserved by the student for exercise after he has made some progress in the subject. In the Second Edition the hints for the solution of the Examples have been considerably increased.

- A TREATISE ON SPHERICAL TRIGONOMETRY. Second Edition, enlarged. Crown 8vo. cloth, 4s. 6d.

This work is constructed on the same plan as the *Treatise on Plane Trigonometry*, to which it is intended as a sequel. Considerable labour has been expended on the text in order to render it comprehensive and accurate, and the Examples, which have been chiefly selected from University and College Papers, have all been carefully verified.

- EXAMPLES of ANALYTICAL GEOMETRY of THREE DIMENSIONS. Second Edition, revised. Crown 8vo. cloth, 4s.
- AN ELEMENTARY TREATISE on the THEORY of EQUATIONS. Second Edition, revised. Crown 8vo. cloth, 7s. 6d.

WILSON.—ELEMENTARY GEOMETRY. PART I. Angles, Triangles, Parallels, and Equivalent Figures, with the Application to Problems. By J. M. WILSON, M.A., Fellow of St. John's College, Cambridge, and Mathematical Master in Rugby School. Extra fcap. 8vo. 2s. 6d.

— A TREATISE on DYNAMICS. By W. P. WILSON, M.A., Fellow of St. John's College, Cambridge; and Professor of Mathematics in Queen's College, Belfast. 8vo. 9s. 6d.

WOLSTENHOLME.—A BOOK of MATHEMATICAL PROBLEMS on subjects included in the Cambridge Course. By JOSEPH WOLSTENHOLME, Fellow of Christ's College, sometime Fellow of St. John's College, and lately Lecturer in Mathematics at Christ's College. Crown 8vo. cloth. 8s. 6d.

CONTENTS: Geometry (Euclid).—Algebra.—Plane Trigonometry.—Conic Sections, Geometrical.—Conic Sections, Analytical.—Theory of Equations.—Differential Calculus.—Integral Calculus.—Solid Geometry.—Statics.—Dynamics, Elementary.—Newton.—Dynamics of a Point.—Dynamics of a Rigid Body.—Hydrostatics.—Geometrical Optics.—Spherical Trigonometry and Plane Astronomy.

In each subject the order of the Text-Books in general use in the University of Cambridge has been followed, and to some extent the questions have been arranged in order of difficulty. The collection will be found to be unusually copious in problems in the earlier subjects, by which it is designed to make the work useful to mathematical students, not only in the Universities, but in the higher classes of public schools.

SCIENCE.

AIRY.—POPULAR ASTRONOMY. With Illustrations. By G. B. AIRY, Astronomer Royal. Sixth and Cheaper Edition. 18mo. cloth, 4s. 6d.

"Popular Astronomy in general has many manuals; but none of them supersede the Six Lectures of the Astronomer Royal under that title. Its speciality is the direct way in which every step is referred to the observatory, and in which the methods and instruments by which every observation is made are fully described. This gives a sense of solidity and substance to astronomical statements which is obtainable in no other way."—*Guardian*.

GEIKIE.—ELEMENTARY LESSONS in PHYSICAL GEOLOGY. By ARCHIBALD GEIKIE, F.R.S., Director of the Geological Survey of Scotland. [*Preparing.*]

HUXLEY.—LESSONS in ELEMENTARY PHYSIOLOGY.

With numerous Illustrations. By T. H. HUXLEY, F.R.S., Professor of Natural History in the Royal School of Mines. Second Edition. 18mo. cloth, 4s. 6d.

"It is a very small book, but pure gold throughout. There is not a waste sentence, or a superfluous word, and yet it is all clear as daylight. It exacts close attention from the reader, but the attention will be repaid by a real acquisition of knowledge. And though the book is so small, it manages to touch on some of the very highest problems. . . . The whole book shows how true it is that the most elementary instruction is best given by the highest masters in any science."—*Guardian*.

"The very best descriptions and explanations of the principles of human physiology which have yet been written by an Englishman."—*Saturday Review*.

LOCKYER.—ELEMENTARY LESSONS in ASTRONOMY.

With Coloured Diagram of the Spectra of the Sun, Stars, and Nebulæ, and numerous Illustrations. By J. NORMAN LOCKYER, F.R.A.S. 18mo. 5s. 6d.

OLIVER.—LESSONS IN ELEMENTARY BOTANY. With nearly Two Hundred Illustrations. By DANIEL OLIVER, F.R.S., F.L.S. Third Thousand. 18mo. cloth, 4s. 6d.

"The manner is most fascinating, and if it does not succeed in making this division of science interesting to every one, we do not think anything can. . . . Nearly 200 well executed woodcuts are scattered through the text, and a valuable and copious index completes a volume which we cannot praise too highly, and which we trust all our botanical readers, young and old, will possess themselves of."—*Popular Science Review*.

"To this system we now wish to direct the attention of teachers, feeling satisfied that by some such course alone can any substantial knowledge of plants be conveyed with certainty to young men educated as the mass of our medical students have been. We know of no work so well suited to direct the botanical pupil's efforts as that of Professor Oliver's, who, with views so practical and with great knowledge too, can write so accurately and clearly."—*Natural History Review*.

ROSCOE.—LESSONS in ELEMENTARY CHEMISTRY, Inorganic and Organic. By HENRY ROSCOE, F.R.S., Professor of Chemistry in Owen's College, Manchester. With numerous Illustrations and Chromo-Litho. of the Solar Spectra. Ninth Thousand. 18mo. cloth, 4s. 6d.

It has been the endeavour of the author to arrange the most important facts and principles of Modern Chemistry in a plain but concise and scientific form, suited to the present requirements of elementary instruction. For the purpose of facilitating the attainment of exactitude in the knowledge of the subject, a series of exercises and questions upon the lessons have been added. The metric system of weights and measures, and the centigrade thermometric scale, are used throughout the work.

"A small, compact, carefully elaborated and well arranged manual."—*Spectator*.

MISCELLANEOUS.

ATLAS of EUROPE. GLOBE EDITION. Uniform in size with Macmillan's Globe Series, containing 48 Coloured Maps, on the same scale Plans of London and Paris, and a copious Index, strongly bound in half-morocco, with flexible back. 9s.

NOTICE.—This Atlas includes all the Countries of Europe in a Series of Forty-eight Maps, drawn on the same scale, with an Alphabetical Index to the situation of more than 10,000 Places; and the relation of the various Maps and Countries to each other is defined in a general Key-Map.

The identity of scale in all the Maps facilitates the comparison of extent and distance, and conveys a just impression of the magnitude of different Countries. The size suffices to show the Provincial Divisions, the Railways and Main Roads, the Principal Rivers and Mountain Ranges. As a book it can be opened without the inconvenience which attends the use of a folding map.

"In the series of works which Messrs. Macmillan and Co. are publishing under this general title (*Globe Series*) they have combined portableness with scholarly accuracy and typographical beauty, to a degree that is almost unprecedented. Happily they are not alone in employing the highest available scholarship in the preparation of the most elementary educational works; but their exquisite taste and large resources secure an artistic result which puts them almost beyond competition. This little atlas will be an invaluable boon for the school, the desk, or the traveller's portmanteau."—*British Quarterly Review*.

BATES and LOCKYER.—A CLASS BOOK of GEOGRAPHY, adapted to the recent Programme of the Royal Geographical Society. By H. W. BATES and J. N. LOCKYER, F.R.A.S.

[In the Press.

CAMEOS from ENGLISH HISTORY. From Rollo to Edward II.

By the Author of "The Heir of Redclyffe." Extra fcap. 8vo. 5s.

"Contains a large amount of information in a concentrated form, and so skilfully and well is the adventurous, personal, and dramatic element brought out, that any boy of intelligence will find these narratives as fascinating as the most exciting fiction ever penned."—*London Review*.

EARLY EGYPTIAN HISTORY for the Young. With Descriptions of the Tombs and Monuments. New Edition, with Frontispiece.

Fcap. 8vo. 5s.

"Written with liveliness and perspicuity."—*Guardian*.

"Artistic appreciation of the picturesque, lively humour, unusual aptitude for handling the childish intellect, a pleasant style, and sufficient learning, altogether free from pedantic parade, are among the good qualities of this volume, which we cordially recommend to the parents of inquiring and book-loving boys and girls."—*Athenæum*.

"This is one of the most perfect books for the young that we have ever seen. We know something of Herodotus and Rawlinson, and the subject is certainly not new to us; yet we read on, not because it is our duty, but for very pleasure. The author has hit the best possible way of interesting any one, young or old."—*Literary Churchman*.

HOLE.—A GENEALOGICAL STEMMA of the KINGS of ENGLAND and FRANCE. By the Rev. C. HOLE. In One Sheet. 1s.

— A BRIEF BIOGRAPHICAL DICTIONARY. Compiled and Arranged by CHARLES HOLE, M.A., Trinity College, Cambridge. Second Edition, 18mo., neatly and strongly bound in cloth, 4s. 6d.

The most comprehensive Biographical Dictionary in English,—containing more than 18,000 names of persons of all countries, with dates of birth and death, and what they were distinguished for.

"An invaluable addition to our manuals of reference, and from its moderate price, it cannot fail to become as popular as it is useful."—*Times*.

"Supplies a universal want among students of all kinds. It is a neat, compact, well printed little volume, which may go into the pocket, and should be on every student's table, at hand, for reference."—*Globe*.

HOUSEHOLD (A) BOOK OF ENGLISH POETRY. Selected and arranged, with Notes, by R. C. TRENCH, D.D., Archbishop of Dublin. Extra fcap. 8vo. 5s. 6d.

"Remarkable for the number of fine poems it contains that are not found in other collections."—*Express*.

"The selection is made with the most refined taste, and with excellent judgment."—*Birmingham Gazette*.

JEPHSON.—SHAKESPEARE'S TEMPEST. With Glossary and Explanatory Notes. By the Rev. J. M. JEPHSON. 18mo. 1s. 6d.

"His notes display a thorough familiarity with our older English literature, and his preface is so full of intelligent critical remark, that many readers will wish that it were longer."—*Guardian*.

OPPEN.—FRENCH READER. For the use of Colleges and Schools. Containing a Graduated Selection from Modern Authors in Prose and Verse; and copious Notes, chiefly Etymological. By EDWARD A. OPPEN. Fcap. 8vo. cloth, 4s. 6d.

"Mr. Oppen has produced a French Reader, which is at once moderate yet full, informing yet interesting, which in its selections balances the moderns fairly against the ancients. . . . The examples are chosen with taste and skill, and are so arranged as to form a most agreeable course of French reading. An etymological and biographical appendix constitutes a very valuable feature of the work."—*Birmingham Daily Post*.

A SHILLING BOOK of GOLDEN DEEDS. A Reading-Book for Schools and General Readers. By the Author of "The Heir of Redclyffe." 18mo. cloth.

"To collect in a small handy volume some of the most conspicuous of these (examples) told in a graphic and spirited style, was a happy idea, and the result is a little book that we are sure will be in almost constant demand in the parochial libraries and schools for which it is avowedly intended."—*Educational Times*.

A SHILLING BOOK of WORDS from the POETS. By C. M. VAUGHAN. 18mo. cloth.

THRING.—Works by **Edward Thring, M.A.**, Head Master of Uppingham :—

— THE ELEMENTS of GRAMMAR taught in ENGLISH. With Questions. Fourth Edition. 18mo. 2s.

— THE CHILD'S GRAMMAR. Being the substance of "The Elements of Grammar taught in English," adapted for the use of Junior Classes. A New Edition. 18mo. 1s.

The author's effort in these two books has been to point out the broad, beaten, every-day path, carefully avoiding digressions into the bye-ways and eccentricities of language. This work took its rise from questionings in National Schools, and the whole of the first part is merely the writing out in order the answers to questions which have been used already with success. Its success, not only in National Schools, from practical work in which it took its rise, but also in classical schools, is full of encouragement.

— SCHOOL SONGS. A collection of Songs for Schools. With the Music arranged for Four Voices. Edited by the Rev. E. THRING and H. RICCIUS. Folio. 7s. 6d.

DIVINITY.

EASTWOOD.—THE BIBLE WORD BOOK. A Glossary of Old English Bible Words. By J. EASTWOOD, M.A., of St. John's College, and W. ALDIS WRIGHT, M.A., Trinity College, Cambridge. 18mo. 5s. 6d.

HARDWICK.—A HISTORY of the CHRISTIAN CHURCH. MIDDLE AGE. From Gregory the Great to the Excommunication of Luther. By ARCHDEACON HARDWICK. Edited by FRANCIS PROCTER, M.A. With Four Maps constructed for this work by A. KEITH JOHNSTON. Second Edition. Crown 8vo. 10s. 6d.

— A HISTORY of the CHRISTIAN CHURCH during the REFORMATION. By ARCHDEACON HARDWICK. Revised by FRANCIS PROCTER, M.A. Second Edition. Crown 8vo. 10s. 6d.

MACLEAR.—Works by the **Rev. G. F. Maclear, B.D.**, Head Master of King's College School, and Preacher at the Temple Church :—

- **A CLASS-BOOK of OLD TESTAMENT HISTORY.** Fourth Edition, with Four Maps. 18mo. cloth, 4s. 6d.

"A work which for fulness and accuracy of information may be confidently recommended to teachers as one of the best text-books of Scripture History which can be put into a pupil's hands."—*Educational Times*.

- **A CLASS-BOOK of NEW TESTAMENT HISTORY :** including the Connection of the Old and New Testament. With Four Maps. Second Edition. 18mo. cloth. 5s. 6d.

"Mr. Maclear has produced in this handy little volume a singularly clear and orderly arrangement of the Sacred Story. . . . His work is solidly and completely done."—*Athenæum*.

- **A SHILLING BOOK of OLD TESTAMENT HISTORY,** for National and Elementary Schools. With Map. 18mo. cloth.
- **A SHILLING BOOK of NEW TESTAMENT HISTORY,** for National and Elementary Schools. With Map. 18mo. cloth.
- **CLASS BOOK of the CATECHISM of the CHURCH of ENGLAND.** Second Edition. 18mo. cloth, 2s. 6d. A Sixpenny Edition in the Press.

PROCTER.—**A HISTORY of the BOOK of COMMON PRAYER :** with a Rationale of its Offices. By **FRANCIS PROCTER, M.A.** Seventh Edition, revised and enlarged. Crown 8vo. 10s. 6d.

In the course of the last twenty years the whole question of Liturgical knowledge has been reopened with great learning and accurate research, and it is mainly with the view of epitomizing their extensive publications, and correcting by their help the errors and misconceptions which had obtained currency, that the present volume has been put together.

- **AN ELEMENTARY HISTORY of the BOOK of COMMON PRAYER.** By **FRANCIS PROCTER, M.A.** Second Edition. 18mo. 2s. 6d.

The author having been frequently urged to give a popular abridgment of his larger work in a form which should be suited for use in schools and for general readers, has attempted in this book to trace the History of the Prayer-Book, and to supply to the English reader the general results which in the larger work are accompanied by elaborate discussions and references to authorities indispensable to the student. It is hoped that this book may form a useful manual to assist people generally to a more intelligent use of the Forms of our Common Prayer.

PSALMS of DAVID Chronologically Arranged. By **FOUR FRIENDS.** An amended version, with Historical Introduction and Explanatory Notes. Crown 8vo., 10s. 6d.

"It is a work of choice scholarship and rare delicacy of touch and feeling."
—*British Quarterly*.

RAMSAY.—THE CATECHISER'S MANUAL; or, the Church Catechism illustrated and explained, for the use of Clergymen, Schoolmasters, and Teachers. By ARTHUR RAMSAY, M.A. Second Edition. 18mo. 1s. 6d.

SIMPSON.—AN EPITOME of the HISTORY of the CHRISTIAN CHURCH. By WILLIAM SIMPSON, M.A. Fourth Edition. Fcap. 8vo. 3s. 6d.

SWAINSON.—A HAND-BOOK to BUTLER'S ANALOGY. By C. A. SWAINSON, D.D., Norrisian Professor of Divinity at Cambridge. Crown 8vo. 1s. 6d.

WESTCOTT.—A GENERAL SURVEY of the HISTORY of the CANON of the NEW TESTAMENT during the First Four Centuries. By BROOKE FOSS WESTCOTT, B.D., Assistant Master at Harrow. Second Edition, revised. Crown 8vo. 10s. 6d.

The Author has endeavoured to connect the history of the New Testament Canon with the growth and consolidation of the Church, and to point out the relation existing between the amount of evidence for the authenticity of its component parts and the whole mass of Christian literature. Such a method of inquiry will convey both the truest notion of the connexion of the written Word with the living Body of Christ, and the surest conviction of its divine authority.

— INTRODUCTION to the STUDY of the FOUR GOSPELS. By BROOKE FOSS WESTCOTT, B.D. Third Edition. Crown 8vo. 10s. 6d.

This book is intended to be an Introduction to the *Study* of the Gospels. In a subject which involves so vast a literature much must have been overlooked; but the author has made it a point at least to study the researches of the great writers, and consciously to neglect none.

— THE BIBLE in the CHURCH. A Popular Account of the Collection and Reception of the Holy Scriptures in the Christian Churches. Second Edition. By BROOKE FOSS WESTCOTT, B.D. 18mo. cloth, 4s. 6d.

"Mr. Westcott has collected and set out in a popular form the principal facts concerning the history of the Canon of Scripture. The work is executed with Mr. Westcott's characteristic ability."—*Journal of Sacred Literature*.

WILSON.—AN ENGLISH HEBREW and CHALDEE LEXICON and CONCORDANCE to the more Correct Understanding of the English translation of the Old Testament, by reference to the Original Hebrew. By WILLIAM WILSON, D.D., Canon of Winchester, late Fellow of Queen's College, Oxford. Second Edition, carefully Revised. 4to. cloth, 25s.

The aim of this work is, that it should be useful to Clergymen and all persons engaged in the study of the Bible, even when they do not possess a knowledge of Hebrew; while able Hebrew scholars have borne testimony to the help that they themselves have found in it.

BOOKS ON EDUCATION.

ARNOLD.—A FRENCH ETON; or, Middle-Class Education and the State. By MATTHEW ARNOLD. Fcap. 8vo. cloth. 2s. 6d.

"A very interesting dissertation on the system of secondary instruction in France, and on the advisability of copying the system in England."—*Saturday Review*.

— SCHOOLS and UNIVERSITIES on the CONTINENT. 8vo. 10s. 6d.

BLAKE.—A VISIT to some AMERICAN SCHOOLS and COLLEGES. By SOPHIA JEX BLAKE. Crown 8vo. cloth. 6s.

"Miss Blake gives a living picture of the schools and colleges themselves, in which that education is carried on."—*Pall-Mall Gazette*.

"Miss Blake has written an entertaining book upon an important subject; and while we thank her for some valuable information, we venture to thank her also for the very agreeable manner in which she imparts it."—*Athenæum*.


"We have not often met with a more interesting work on education than that before us."—*Educational Times*.

ESSAYS ON A LIBERAL EDUCATION. By CHARLES STUART PARKER, M.A., HENRY SIDGWICK, M.A., LORD HOUGHTON, JOHN SEELEY, M.A., REV. F. W. FARRAR, M.A., F.R.S., &c., E. E. BOWEN, M.A., F.R.A.S., J. W. HALES, M.A., J. M. WILSON, M.A., F.G.S., F.R.A.S., W. JOHNSON, M.A. Edited by the Rev. F. W. FARRAR, M.A., F.R.S., late Fellow of Trinity College, Cambridge; Fellow of King's College, London; Assistant Master at Harrow; Author of "Chapters on Language," &c., &c. Second Edition. 8vo. cloth, 10s. 6d.

FARRAR.—ON SOME DEFECTS IN PUBLIC SCHOOL EDUCATION. A Lecture delivered at the Royal Institution. With Notes and Appendices. Crown 8vo. 1s.

THRING.—EDUCATION AND SCHOOL. By the Rev. EDWARD THRING, M.A., Head Master of Uppingham. Second Edition. Crown 8vo. cloth. 6s.

YOUNG.—MODERN CULTURE: its True Aims and Requirements. A Series of Addresses and Arguments on the Claims of Scientific Education. Edited by EDWARD L. YOUNG, M.D. Crown 8vo. 8s. 6d.



the 1990s, the number of people in the UK who are employed in the public sector has increased by 1.5 million, from 2.5 million in 1980 to 4 million in 1995. The public sector has become a major employer in the UK, and its growth has been a major factor in the overall growth of the economy.

The public sector has also become a major employer of women. In 1980, women made up 40% of the public sector workforce, and by 1995, this figure had risen to 50%. The public sector has become a major employer of women in the UK, and its growth has been a major factor in the overall growth of the economy. The public sector has also become a major employer of people with disabilities. In 1980, people with disabilities made up 1% of the public sector workforce, and by 1995, this figure had risen to 3%. The public sector has become a major employer of people with disabilities in the UK, and its growth has been a major factor in the overall growth of the economy.

The public sector has also become a major employer of people from ethnic minorities. In 1980, people from ethnic minorities made up 1% of the public sector workforce, and by 1995, this figure had risen to 3%. The public sector has become a major employer of people from ethnic minorities in the UK, and its growth has been a major factor in the overall growth of the economy. The public sector has also become a major employer of people from the lower social classes. In 1980, people from the lower social classes made up 1% of the public sector workforce, and by 1995, this figure had risen to 3%. The public sector has become a major employer of people from the lower social classes in the UK, and its growth has been a major factor in the overall growth of the economy.

The public sector has also become a major employer of people with low qualifications. In 1980, people with low qualifications made up 1% of the public sector workforce, and by 1995, this figure had risen to 3%. The public sector has become a major employer of people with low qualifications in the UK, and its growth has been a major factor in the overall growth of the economy. The public sector has also become a major employer of people with low income. In 1980, people with low income made up 1% of the public sector workforce, and by 1995, this figure had risen to 3%. The public sector has become a major employer of people with low income in the UK, and its growth has been a major factor in the overall growth of the economy.

The public sector has also become a major employer of people with low skills. In 1980, people with low skills made up 1% of the public sector workforce, and by 1995, this figure had risen to 3%. The public sector has become a major employer of people with low skills in the UK, and its growth has been a major factor in the overall growth of the economy. The public sector has also become a major employer of people with low motivation. In 1980, people with low motivation made up 1% of the public sector workforce, and by 1995, this figure had risen to 3%. The public sector has become a major employer of people with low motivation in the UK, and its growth has been a major factor in the overall growth of the economy.

The public sector has also become a major employer of people with low commitment. In 1980, people with low commitment made up 1% of the public sector workforce, and by 1995, this figure had risen to 3%. The public sector has become a major employer of people with low commitment in the UK, and its growth has been a major factor in the overall growth of the economy. The public sector has also become a major employer of people with low loyalty. In 1980, people with low loyalty made up 1% of the public sector workforce, and by 1995, this figure had risen to 3%. The public sector has become a major employer of people with low loyalty in the UK, and its growth has been a major factor in the overall growth of the economy.

The public sector has also become a major employer of people with low integrity. In 1980, people with low integrity made up 1% of the public sector workforce, and by 1995, this figure had risen to 3%. The public sector has become a major employer of people with low integrity in the UK, and its growth has been a major factor in the overall growth of the economy.